CUMULATIVE AUTHOR INDEX
Environmental Entomology
Volume 37, 2008

A

Abbasipour, H., 38
Abel, C. A., 1538
Adair, R. C., Jr, 914
Adamczyk, J. J., 1396
Addicott, J. F., 1017
Allender, C. J., 817
Alleyne, M., 1307
Alston, D. G., 1272
Alvarez, J. M., 592
Amarasekare, K. G., 630
Andersen, P. C., 1049
Anulewicz, A. C., 230
Arbab, A., 1381
Arbogast, R. T., 131
Ayres, M. P., 650

B

Baba, Y. G., 938
Babu, A., 1416
Bahlai, C. A., 973
Bai, B. B., 277
Barbehenn, R. V., 1113
Baselga, A., 1488
Bauer, L. S., 1344
Bean, C., 897
Beaver, R. A., 1461
Behan-Pelletier, V. M., 464
Behle, R. W., 162, 1126
Benbow, M. E., 1162
Bennett, G. W., 741
Berenbaum, M. R., 601
Berkebile, D. R., 625
Beroiz, B., 1354
Billings, R. F., 650
Blackburn, L. M., 87
Blanco, C. A., 1538
Bloem, K. A., 1291
Bloem, S., 1291
Boano, G., 1145
Boettner, G., 1429
Boiteau, G., 575
Boivin, G., 907
Boone, C. K., 150
Bosque-Pérez, N. A., 592
Bourassa, S., 1214
Bower, N. W., 871
Bowling, D., 1373
Boyero, J. R., 1531
Brady, J. A., 1525
Branson, D. H., 686
Brent, C. S., 883
Broatch, J. S., 1451
Brodbeck, B. V., 1049
Brown, R. L., 453
Brunner, J. F., 192, 679
Buczkowski, G., 741
Burd, J. D., 586
Butler, C. D., 964

C

Capinera, J. L., 1565
Cappaert, D. L., 230
Cárcamo, H. A., 1214
Cardoza, Y. J., 956
Carel, Y., 307
Carpenter, J. E., 1291
Carrière, Y., 1514
Carrillo, D., 1565
Carroll, A. S., 586
Castañera, P., 1354
Cazares, C. L., 876
Chamberlain, S. A., 808
Chapman, P. S., 1086
Chen, X.-X., 1264
Cheng, D.-F., 774
Cheng, J., 748
Childers, C. C., 555
Chong, J.-H., 323
Clancy, K. M., 94, 817
Clark, T. L., 247
Clayton, G. W., 1451
Corlay, F., 907
Corn, J. G., 510
Corronca, J. A., 419
Côté, M.-J., 636
Cottrell, T. E., 162, 1508
Coubes, M., 307
Crist, T. O., 897
Cristofaro, M., 1185
Crook, D. J., 356
Crowder, D. W., 1514
Cruz-López, L., 876
Cuda, J. P., 862

D

Dalthorp, D., 409
Damborsky, M. P., 419
Dashiell, K. E., 247
David, J. F., 45
Davis, A.L.V., 431
Decker, K. B., 670
Degomez, T. E., 57, 94, 817
Delaney, K. J., 1332
De La Poza, M., 1354
DeLucia, E. H., 601
Descampe, A., 224
Desurmont, G. A., 520, 845, 1241
Diaz, M.D.C., 520, 1438
Diaz, R., 862
Dickenson, R. M., 782
Dively, G. P., 1
Dixon, P. M., 1
Doblas-Miranda, E., 857
Dole, S. A., 1461
Dong, L., 263
Dorhout, D. L., 1280
Dorn, S., 1291
Dosdall, L. M., 1451
Dowd, M. K., 1081
Dreves, A. J., 409
Du, L., 29
Duan, J. J., 1, 278, 838
Duke, S. E., 1081
Dunlap, C. A., 1126
Dupont, M. M., 1224
Duyck, P. F., 307
Dyer, L. A., 947

E

Eigenbrode, S. D., 592
Elkinton, J. S., 1429
Ellersieck, M. R., 1558
Elliott, N. C., 315, 1063
Ellsworth, P. C., 1514
Embrey, S. K., 255
Emerick, J. J., 871
Epsky, N. D., 630, 1119
Esiekpe, O., 478

F

Fadamiro, H. Y., 555
Farinós, G. P., 1354
Fathipour, Y., 38
Feng, H.-Q., 774
Feng, M.-G., 1264
Fidgen, J. G., 1498
Fiedler, A. K., 707
Fisher, J., 1126
Fletcher, M. R., 546
Floris, I., 505
Francese, J. A., 356
Frank, S. D., 478
Fraser, I., 356
Freeland, T. B., Jr, 198
French, B. W., 247
Friedenberg, N. A., 650
Fuest, J., 1508
Fuxa, J. R., 755

G

Gañán, I., 1488
Gandhi, K.J.K., 725
Gao, F., 29
Gao, X. W., 1410
García, Á. H., 1531

Gardner, W. A., 162
Garthwaite, D., 546
Gaylord, M. L., 57
Ge, F., 29
Ghimire, M. N., 1300
Giles, K. L., 315, 1063
Gilles, J., 45
Goane, L., 1025
Goenaga, R., 110
Golizadeh, A., 38
González-Hernández, H., 876
Grace, J. K., 400
Graf, G., 472
Greenberg, S. M., 1396
Grinberg, P. S., 636
Groves, R. L., 79, 787
Gu, H.-N., 374
Guillemaud, T., 293
Gültekin, L., 1185
Guo, J.-Y., 263, 1035
Guo, Y.-Y., 774, 1410
Gwynne, D. T., 889

H

Hagler, J. R., 787
Hahn, D. A., 1074
Haile, F. J., 1332
Hakkarainen, H., 51
Halbert, S. E., 586
Hall, D. G., 914
Hall, D. R., 719
Hallett, R. H., 973
Hamilton, B., 1099
Hamilton, G. C., 348
Hammons, D. L., 608
Han, L.-Z., 374, 1361
Han, Z.-J., 263
Hance, T., 224
Handelsman, J., 1344
Hanks, L. M., 1307
Hanula, J. L., 1403
Harding, K. M., 546
Hargroder, T., 555
Harker, K. N., 1451
Harwood, J. D., 525, 990
Hawkins, L., 980
Heath, R. R., 1119
Hebbar, P., 719
Hebertson, E. G., 281
Heimpel, G. E., 301
Heist, E. J., 271
Hellmich, R. L., 1, 615
Helms, K. R., 487
Hentz, M. G., 586, 914
Herbert, J. J., 1258
Herman, R. A., 255
Herms, D. A., 242
Hernández-Crespo, P., 1354
Hibbard, B. E., 247, 1558
Higdon, M. L., 1558
Higgins, L. S., 1
Hight, S. D., 1291
Higley, L. G., 1332
Hill, J. G., 453
Hilton, R., 980
Hodge, S., 1573
Hoelmer, K. A., 391
Hoffman Gray, R., 1174
Hofstetter, R. W., 57, 70
Horn, D. J., 725, 1258
Horton, D. L., 1508
Hu, C., 825
Hu, J.-H., 206
Huang, M., 1043
Huang, Z.-H., 1264
Huesing, J. E., 838
Hulcr, J., 1461
Hunter, W. B., 1049
Husseneder, C., 400

I

Isaacs, R., 707

J

James, J., 1373
Jenkins, D. A., 110
Jenkins, M. J., 281
Jiang, C., 838
Jiang, C.-Y., 825
Jiang, H.-B., 1105
Jiang, X.-F., 1389
Joern, A., 333
Johnson, A. V., 1224
Johnson, M. W., 787
Jonas, J. L., 333
Jones, D. B., 1063
Jones, V. P., 192, 679
Joost, P. H., 213

K

Kamali, K., 38
Kang, L., 29
Karumuru, B., 719
Kattes, D., 11
Keathley, C. P., 1548
Keena, M. A., 636
Keim, P., 817
Kendra, P. E., 1119
Kennedy, G. G., 79, 1154
Ketner, K., 980
Khrimian, A., 356
Kim, H., 660
Kim, H. G., 1272
Kim, K. S., 293
King, B. H., 782
Klepzig, K. D., 956
Knight, A. L., 980, 1136
Knoester, J. J., 1113
Knopper, L. D., 546
Koch, F. H., 442
Kohler, G. R., 494, 1477
Kok, L. T., 1498
Kontodimas, D. C., 1381
Koppenhöfer, A. M., 1126
Kouchoukos, N., 650
Kratz, C. J., 734
Krehbiel, R. L., 1224
Krupke, C. H., 192
Kuang, L. F., 719
Kumm, S., 1422
Kurtural, S. K., 608

L

Laird, C., 1126
Laird, R. A., 1017
Lambert, B. D., 11
Lambert, S., 719
Landis, D., 707
Larney, F. J., 1214
Larsen, K. J., 1471
Larsen, T. E., 1136
Larson, T. L., 980
Lauzire, I., 925
Layne, J. R., Jr, 1069
Lecomte, P., 45
Lee, J.-H., 660
Leszczynski, C. F., 1069
Lewis, L. C., 615
Liebman, M., 121
Liere, H., 143
Lindo, Z., 464
Liu, J.-C., 1105
Liu, P., 1361
Liu, Q., 1525
Liu, S.-S., 1264
Liu, T.-X., 1396
Liu, W.-X., 1035
Livingston, G. F., 734
Loaiza, V., 333
Longnecker, M. T., 796
López, J. D., Jr, 1081
López-Collado, J., 876
López-Guillén, G., 876
Lorch, P. D., 889
Lorimer, C. G., 1174
Lou, Y., 748
Lövei, G. L., 263
Luo, L.-Z., 1389
Lv, J., 796
Lynch, D. H., 575
Lyytikäinen-Saarenmaa, P., 1582

M

Maben, R. E., 1113
Macagno, A.L.M., 1145
Macleod, E. C., 973
Maisin, N., 719
Malo, E. A., 876
Mannion, C. M., 323, 630
Manrique, V., 862
Marshall, D. B., 366
Martin, A. R., 1074
Martin, R. C., 575
Martínez-Ferri, E., 1531
Massad, T. J., 947
Mastro, V. C., 356

Matadha, D., 348
Mathur, S., 472
Matsuura, K., 400
Mattson, W. J., 1003
Mausel, D. L., 1498
Mazzocchetti, A., 382
McCormick, R., 255
McCullough, D. G., 230
McMillin, J. D., 57, 94, 817
McNeill, M. R., 1381
Meck, M. C., 1154
Meskens, C., 224
Miller, A., 94
Miller, N. J., 293
Mineau, P., 546
Mironidis, G. K., 16
Mitchell, F., 925
Miyashita, T., 938
Mizell, R. F., III, 1049
Montgomery, W. S., 1119
Morales-Ramos, J. A., 1538
Moreira, G.R.P., 1323
Moritz, G., 1422
Morsello, S. C., 79
Moser, J. C., 956
Moser, S. E., 525, 1307
Moya-Raygoza, G., 1471
Muñoz, S. T., 1531
Muraleedharan, N., 1416
Murphy, S. T., 1035
Myers, H. M., 11

N

Nadel, H., 787
Nagata, R. T., 586
Nansen, C., 1373
Nault, B. A., 79
Nay, J. E., 568
Nesbitt, M., 555
Nielsen, A. L., 348
Northfield, T., 1049
Novoa, F., 1488
Novy, R. G., 592
Nuessly, G. S., 586

O

Obrycki, J. J., 525
Oishi, M., 1231
O'Neil, R. J., 964
O'Neill, B. F., 601
O'Rourke, M. E., 121
Ortego, F., 1354
Osborne, L. S., 630
Overholt, W. A., 862
Oyediran, I. O., 247

P

Palestrini, C., 1145
Papadopoulos, G. D., 340
Papadoulis, G. T., 340
Parajulee, M. N., 29,
Parry, D., 1429
Pasonen, H.-L., 1582
Pasteels, J., 224
Pearson, C. V., 947
Peck, D. C., 1438
Peña, J. E., 1565
Peng, Y.-F., 825, 1361
Pérez, M. J., 1531
Perfecto, I., 143, 181
Perring, T. M., 568
Peterson, R.K.D., 1332
Pfannenstiel, R. S., 996
Pfeiffer, D. G., 1208
Phillips, T. W., 1300
Philpott, S. M., 181
Phoofolo, M. W., 315
Pike, K. S., 472
Polak, M., 1099
Poland, T. M., 230, 356
Porter, P., 1373
Porter, S. D., 1074
Portilla, M., 1538
Potter, D. A., 608, 1548
Powell, G., 1573
Powell, J. E., 400
Prasifka, J. R., 1
Preisser, E., 1429
Puckhaber, L. S., 1081
Puranasakul, W., 1461
Pureswaran, D. S., 70
Purrington, F. F., 725
Purung, H. B., 719

R

Raffa, K. F., 150, 956, 1174, 1344
Rantala, M. J., 51
Rashed, A., 1099
Raworth, D. A., 472
Rebek, E. J., 242
Reeve, J. D., 271
Ren, N., 748
Rendón, P. A., 764
Reyes-López, J., 857
Rice, M. E., 121, 1280
Richter, A., 755
Riley, D. G., 213
Rimoldi, F., 1252
Ring, D., 755
Robertson, I. C., 850
Roda, A. L., 323
Rodrigues, D., 1323
Roitberg, B. D., 534
Rojas, J. C., 876
Rojas, M. G., 1538
Rolando, A., 1145
Ronco, A. E., 1252
Ross, D. W., 494, 1477
Rubio, G. D., 419
Ruiu, L., 505
Runkle, J. R., 696

S

Sachin, J. P., 1416
Salom, S. M., 1498
Sanford, M. R., 1525
Sappington, T. W., 293, 1280, 1396
Sarkar, S., 650
Sarvary, M. A., 1291
Sato, H., 1231
Satta, A., 505
Savopoulou-Soultani, M., 16
Sawyer, A. J., 356
Schaafsma, A. W., 973
Schlising, R. A., 808
Schloss, P. D., 1344
Schneider, M. I., 1252
Scholtz, C. H., 431
Schrey, A. W., 271
Schrey, N. M., 271
Schweikert, Y. M., 1558
Sciarretta, A., 382
Sears, M. K., 973
Seligmann, R., 787
Selvasundaram, R., 1416
Setamou, M., 1396
Seto, M., 1438
Shan, G., 255
Shapiro-Ilan, D. I., 162
Sheather, S., 925
Shrewsbury, P. M., 478
Silverman, B., 725
Simmons, A. M., 391
Sing, S. E., 131
Sisterson, M. S., 1200
Sivinski, J., 764
Six, D. L., 150
Smith, L., 1185
Smith, Lincoln, 510
Smith, W. D., 442
Smitley, D. R., 242
Snodgrass, G. L., 1538
Snyder, A. I., 871
Snyder, M. A., 871
Soares, G.L.G., 1323
Sonthichai, S., 1461
Sorvari, J., 51
Spence, J. R., 1214
Srinivasan, R., 592
Stasek, D. J., 897
Stassi, M., 1145
Stenger, D. C., 787
Stiefel, V. L., 494, 1477
Stipanovic, R. D., 1081
Stolz, U., 293
Story, J. M., 510
Sullivan, B. T., 70, 1403
Sumera, A., 748
Sumerford, D. V., 293
Summerville, K. S., 453, 1224
Sun, J.-Z., 755
Sun, Y.-C., 29
Sword, G. A., 889

T
Tabashnik, B. E., 1514
Takada, M., 938
Tanigoshi, L. K., 472
Tate, C. D., 615
Taylor, D. B., 625
Taylor, R. M., 996
Teixeira, D., 838
Terada, S., 938
Terán-Vargas, A. P., 1538
Tillard, E., 45
Tipping, C., 1049
Tobin, P. C., 87, 1174
Toledo, J., 876
Tomberlin, J. K., 11, 1525
Traniello, J.F.A., 883
Trematerra, P., 382
Trier, T. M., 1003
Trimble, R. M., 366
Tronci, C., 1185
Tshikae, B. P., 431
Tuell, J. K., 707

U
Umeh, V., 555

V
Valladares, G., 1025
Vandermeer, J., 181
van Herk, W. G., 534
Vargo, E. L., 400, 883
Vasanthakumar, A., 1344
Vayssières, J. F., 307
Vela, J. M., 1531
Vernon, R. S., 534
Vihervuori, L., 1582
Vinson, S. B., 487
Virdiana, I., 719
Vittum, P. J., 1126
Vogt, J. T., 198

W
Wagner, D. M., 817
Wagner, M. R., 57
Waits, E. R., 293
Walgenbach, J. F., 1154
Wallet, B., 198
Wallin, K. F., 494, 1477
Wallner, W. E., 636
Wan, F.-H., 263, 1035
Wan, P., 1043
Wang, F., 1361
Wang, J.-J., 1105
Wang, L., 301
Wang, S., 719
Wang, X., 748
Wang, Z.-Y., 1105
Weber, D. C., 1371
Welsman, J. A., 973
Weston, P. A., 520, 845, 1241
White, A. M., 734
White, L. J., 510
Whiteside, M., 546
Wiedenmann, R. N., 1307
Williams, K. K., 57, 94
Williams, M. R., 1313
Williams, W. I., 850
Willink, E., 1025
Wilson, J., 696
Wilson, L. T., 796
Wiman, N. G., 679
Winchester, N. N., 464
Windsor, D., 224
Wood, B. W., 162
Woods, J. N., 696
Woolbright, S. A., 817
Wu, B., 774
Wu, J., 1043
Wu, J.-C., 206
Wu, K., 301, 774, 1043, 1361, 1410
Wu, X.-F., 774
Wyckhuys, K.A.G., 301

X
Xiang, C., 748
Xiao, Y., 555
Xu, H.-W., 206

Y
Yanagi, Y., 938
Yang, G.-Q., 206
Yang, R.-C., 1451
Yao, H.-W., 825
Ye, G.-Y., 825
Yeargan, K. V., 670
Yee, W. L., 172, 1086
Yin, J.-L., 206
Yokoyama, V. Y., 764
Yu, D., 1043
Yu, H., 1410
Yu, Z., 1525

Z
Zangerl, A. R., 601
Zhai, B.-P., 374
Zhang, A., 719
Zhang, G.-F., 1035
Zhang, L., 1389
Zhang, X., 1208
Zhang, X.-X., 374
Zhang, Ying, 301
Zhang, Yongjun, 1410
Zhao, X.-C., 774
Zheng, Y., 150
Zhu, S.-R., 29
Zinni, A., 382

CUMULATIVE KEY WORD INDEX
Environmental Entomology
Volume 37, 2008

A

α diversity,
β diversity, 1461
dipterocarp forest, 1461
effective number of species, 1461
ethanol trap, 1461
flight intercept trap, 1461
abundance,
Dendroctonus, 94
flight seasonality, 94
Ips, 94
Pinus ponderosa, 94
acalypha,
development, 630
hibiscus, 630
parthenium, 630
plumeria, 630
Acanthoscelides obtectus,
biological control, 131
Callosobruchus spp., 131
stored products, 131
Zabrotes subfasciatus, 131
Adelges tsugae, 494, 1477, 1498
Chamaemyiidae, 1477
classical biological control, 1498
exclusion cage, 1498
field insectary, 1498
Laricobius nigrinus, 494
Leucopis argenticollis, 494
Leucopis atrifacies, 494
parasitoids, 1477
predators, 1477
synchrony, 1498
Tsuga heterophylla, 494, 1477
adult feeding,
blood, 625
Stomoxys, 625
sugar, 625
adult life history,
Dacini, 307
host–plant effect, 307
preimaginal development, 307
temperature, 307
aerobic metabolism,
cryobiology, 1069
freeze tolerance, 1069
glycerol, 1069
Hypercompe scribonia, 1069
aerobiology,
biological invasions, 87
local indicator of spatial autocorrelation, 87
Lymantria dispar, 87
quantile regression, 87
African savannas,
bait types, 431
Botswana, 431
Scarabaeinae, 431
trophic associations, 431
Ageniaspis citricola,
Cirrospilus neotropicus, 1025
citrus leafminer, 1025
host selection, 1025
tritrophic interactions, 1025
Agrilus planipennis,
coevolution, 242
host plant resistance, 242
invasive species, 242
wood borers, 242
Agriotes obscurus,
dusky wireworm, 534
insecticide, 534
repellency, 534
agro-ecology,
beetle communities, 121
biological control, 121
generalist predators, 121
natural enemies, 121
Alatae,
growth rate, 1573
host choice, 1573
plant pathogen, 1573
virus vector, 1573
alfalfa,
arthropods, 947
chemical defense, 947
diversity cascades, 947
saponins, 947
4-allylanisole,
Dendroctonus ponderosae, 871
estragole, 871
mountain pine beetle, 871
Pinus ponderosa, 871
alternate hosts,
Diabrotica barberi, 247
prairie grasses, 247
resistance management, 247
alternate weed hosts,
insect vectors, 592
vector performance, 592
vector reservoir, 592
viral inoculum source, 592
ambrosia beetle,
climate matching, 442
cost-weighted distance, 442
redbay, 442
Xyleborus glabratus, 442
amplified fragment length polymorphism,
D. frontalis, 817
Dendroctonus brevicomis, 817
genetic differentiation, 817
Ips pini, 817
Anabrus simplex,
collective movement, 889
hopper band, 889
katydid, 889
radiotransmitter, 889
Anagrus nilaparvatae,
host suitability, 748
infestation level, 748
Nilaparvata lugens, 748
tritrophic interaction, 748
Anastrepha obliqua, 110, 876
Anastrepha suspensa, 110
calling behavior, 876
Diptera, 876
male volatiles, 876
Mangifera indica, 110
Tephritidae, 876
Utetes anastrephae, 110
Anastrepha suspensa,
Anastrepha obliqua, 110
Mangifera indica, 110
Utetes anastrephae, 110
animal foraging,
daily variations, 857
harvester ants, 857
soil temperature, 857
trail system, 857
annual bluegrass weevil,
Poa annua, 1438
population fluctuations, 1438
turfgrass, 1438
Anomala orientalis,
mating disruption, 1126
sex pheromone, 1126
turfgrass, 1126
(Z) and (E)-7-tetradecen-2-one formulation, 1126
ant,
leafhoppers, 1471
mutualism, 1471
shade, 1471
shelter, 1471
ant attendance,
Coccinellidae, 1258
Monomorium, 1258
soybean aphid, 1258
ant-hemipteran mutualism,
Coccinellidae, 143
coffee, 143

Mexico, 143
myrmecophily, 143
Anthonomus grandis grandis,
boll weevil, 1396
cotton, 1396
photoperiod, 1396
anthrone,
extrafloral, 996
nectar, 996
nutrition, 996
spider, 996
Antonina graminis,
Cynodon dactylon, 487
honeydew, 487
invasive species, 487
Solenopsis invicta, 487
Aphidius ervi,
geographical distribution, 472
phenology, 472
population dynamics, 472
Praon unicum, 472
aphid-pathogenic fungi,
colonization, 1264
dissemination, 1264
host dispersal flight, 1264
parasitoids, 1264
aphids,
phytophage, 1531
resistance, 1531
sprouting, 1531
susceptibility, 1531
Aphis glycines, 301, 973
buckthorn, 973
dispersal, 301
environmental condition, 301
flight ability, 301
flight mill, 301
Harmonia axyridis, 973
preference assay, 973
soybean aphid, 973
Aphis gossypii, 29, 263, 1035
Chrysopa pallens, 263
cotton cultivar, 29
elevated CO_2, 29
nontarget effects, 1035
Orius sauteri, 1035
Propylaea japonica, 29
transgenic *Bt* cotton, 263, 1035
tri-trophic interaction, 29, 263, 1035
Apis mellifera,
honey bee, 808
invasive species, 808
native plant, 808
Triteleia laxa, 808
apple, 980, 1136, 1154
competitive attraction, 1136
Cydia pomonella, 1136
diapause, 1154
low volume, 980
mating disruption, 980, 1136
pear, 980
pest management, 980, 1136
Rhagoletis pomonella, 1154
arbuscular mycorrhizal fungi,
Helianthus annuus, 1017
sunflower beetle, 1017
trait-mediated indirect effects, 1017
Zygogramma exclamationis, 1017
ArcGIS mapping,
biological control, 862
developmental rate, 862
Episimus utilis, 862
Schinus terebinthifolius, 862
Argentina,
contiguous habitats, 419
Humid Chaco ecoregion, 419
Mburucuya National Park, 419
spider diversity, 419
Argentine ant,
competition, 741
Linepithema humile, 741
odorous house ant, 741
Tapinoma sessile, 741
arrhenotoky,
haplodiploidy, 1422
thelytoky, 1422
Thysanoptera, 1422
Wolbachia, 1422
arthropods,
alfalfa, 947
chemical defense, 947
diversity cascades, 947
saponins, 947
attractant,
cotton, 1410
natural enemy, 1410
plant volatile, 1410
Auchenorrhyncha,
glassy-winged sharpshooter, 925
Hexapoda, 925
leafhopper, 925
Pierce's disease, 925
augmentative release,
biological control, 1241
Podisus maculiventris, 1241
predation dynamics, 1241
predator behavior, 1241
autumn migration,
flight behavior, 774
light trapping, 774
Mythimna separata, 774
radar observation, 774

B

β diversity,
α diversity, 1461
dipterocarp forest, 1461
effective number of species, 1461
ethanol trap, 1461
flight intercept trap, 1461
Bacillus thuringiensis, 1, 1354
experimental design, 1
gene flow, 1354
genetically modified, 1
Mediterranean corn borer, 1354
RAPD, 1354
risk assessment, 1
transgenic maize, 1354
Bactrocera oleae (Gmelin),
Bactrocera oleae (Gmelin), 764
larval parasitoid, 764
Olea europaea L., 764
bagworm,
competition, 1307
host discrimination, 1307
host feeding, 1307
parasitoid, 1307
bait types,
African savannas, 431
Botswana, 431
Scarabaeinae, 431
trophic associations, 431
bark beetles,
Histiogaster, 956
Hyphomycetes, 956
Leptographium, 956
symbiosis, 956
bark volatiles,
Buprestidae, 356
7-*epi*-sesquithujene, 356
Manuka oil, 356
Phoebe oil, 356
Beauveria bassiana,
biological control, 162
Curculio caryae, 162
pecan, 162
beetle banks,
conservation biological control, 478
habitat manipulation, 478
insectary plants, 478
native plants, 478
beetle communities,
agro-ecology, 121
biological control, 121
generalist predators, 121
natural enemies, 121
behavior, 1049, 1373
cotton gin trash, 1373
glassy-winged sharpshooter, 1049
management, 1049
seasonal dynamics, 1373
soil moisture preference, 1373
Xylella fastidiosa, 1049
Bemisia tabaci,
insect resistance management, 1514
modeling, 1514

operational and environmental factors, 1514
pyriproxyfen, 1514
Bemisia tabaci biotype B,
biological control, 391
Encarsia spp., 391
Eretmocerus spp., 391
sweetpotato whitefly, 391
beneficial acclimation hypothesis,
calorimetry, 1003
diet switching, 1003
dietary acclimation hypothesis, 1003
nutritional ecology, 1003
beneficial arthropods,
Dialeurodes citri, 555
Lepidosaphes beckii, 555
Panonychus citri, 555
Satsuma mandarin, 555
beta diversity,
Calathus, 1488
completeness of inventories, 1488
Iberian Peninsula, 1488
species richness, 1488
biological control, 121, 131, 162, 391, 734, 862, 1185, 1241
Acanthoscelides obtectus, 131
agro-ecology, 121
ArcGIS mapping, 862
augmentative release, 1241
Beauveria bassiana, 162
beetle communities, 121
Bemisia tabaci biotype B, 391
Callosobruchus spp., 131
Curculio caryae, 162
developmental rate, 862
Encarsia spp., 391
Episimus utilis, 862
Eretmocerus spp., 391
generalist predators, 121
host plant, 1185
indirect competition, 734
life history, 1185
Lixinae, 1185
mutualism, 734
natural enemies, 121
pecan, 162
phenology, 1185
Podisus maculiventris, 1241
predation dynamics, 1241
predator behavior, 1241
scale insects, 734
Schinus terebinthifolius, 862
shade coffee, 734
stored products, 131
sweetpotato whitefly, 391
Zabrotes subfasciatus, 131
biological invasions,
aerobiology, 87
local indicator of spatial autocorrelation, 87
Lymantria dispar, 87
quantile regression, 87
biological management,
Diptera, 505
entomopathogenic bacterium, 505
microbial insecticide, 505
treatments, 505
biotechnology,
Cry3Bb1, 838
nontarget organisms, 838
potassium arsenate, 838
protease inhibitor, 838
Black Belt Prairie,
ecological assembly, 453
Formicidae, 453
Mississippi, 453
Solenopsis, 453
black soldier fly larvae,
dairy manure, 1525
Escherichia coli, 1525
Hermetia illucens, 1525
Stratiomyidae, 1525
blood,
adult feeding, 625
Stomoxys, 625
sugar, 625
body size, 51, 1074, 1074
cooling rate, 1074 1074
cooling rate interaction, 1074, 1074
encapsulation, 51
fire ant, 1074, 1074
habitat quality, 51
immunity, 51
supercooling point, 1074, 1074
trade-off, 51
boll weevil,
Anthonomus grandis grandis, 1396
cotton, 1396
photoperiod, 1396
Bombyx mori,
Bt rice, 825
pathological effect, 825
pollen grains, 825
toxic effect, 825
Bostrichidae,
Curculionidae, 1300
host finding, 1300
learning, 1300
Pteromalidae, 1300
Botswana,
African savannas, 431
bait types, 431
Scarabaeinae, 431
trophic associations, 431
Brassica napus,
Brassica rapa, 1451
Delia platura, 1451
Delia radicum, 1451
logistic model, 1451
Brassica oleracea,
development rate, 38
Diadegma anurum, 38
Plutella xylostella, 38
temperature, 38
Brassica rapa,
Brassica napus, 1451
Delia platura, 1451
Delia radicum, 1451
logistic model, 1451
breeding system,
Isoptera, 400
microsatellite genotyping, 400
morphometry, 400
Rhinotermitidae, 400
Bt accumulation in soil,
Cry1F protein, 255
environmental risk, 255
enzyme-linked immunosorbent assay, 255
transgenic corn, 255
Bt cotton,
common cutworm, 1043
population dynamics, 1043
survival, 1043
Bt rice,
Bombyx mori, 825
pathological effect, 825
pollen grains, 825
toxic effect, 825
buckthorn,
Aphis glycines, 973
Harmonia axyridis, 973
preference assay, 973
soybean aphid, 973
bumble bee, 707, 719
habitat management, 707, 719
pollination, 707, 719
prairie plants, 707, 719
restoration, 707, 719
Buprestidae,
bark volatiles, 356
7-*epi*-sesquithujene, 356
Manuka oil, 356
Phoebe oil, 356
butterflies,
day-flying moths, 1313
estimating species richness, 1313
sample completeness, 1313
sampling fraction, 1313

C

cabbage root fly,
crop rotation, 409
Delia radicum, 409
spatial dynamics, 409
spatial ecology, 409
Cactoblastis cactorum,
dispersal, 1291
flight, 1291

overwintering, 1291
seasonality, 1291
cadaverine,
Caribbean fruit fly, 1119
electroantennography, 1119
putrescine, 1119
calamenene,
α-copaene, 1403
exotic species, 1403
Calathus,
beta diversity, 1488
completeness of inventories, 1488
Iberian Peninsula, 1488
species richness, 1488
calling behavior,
Anastrepha obliqua, 876
Diptera, 876
male volatiles, 876
Tephritidae, 876
Callosobruchus spp.,
Acanthoscelides obtectus, 131
biological control, 131
stored products, 131
Zabrotes subfasciatus, 131
calorimetry,
beneficial acclimation hypothesis, 1003
diet switching, 1003
dietary acclimation hypothesis, 1003
nutritional ecology, 1003
canopy lichens,
distribution patterns, 464
microhabitat specificity, 464
montane forests, 464
Oribatid mites, 464
capture-recapture,
Cormack-Jolly-Seber, 1145
survival, 1145
vagility, 1145
Carabidae,
corridor, 725
edge effects, 725
fragmentation, 725
ground beetles, 725
Caribbean fruit fly,
cadaverine, 1119
electroantennography, 1119
putrescine, 1119
Cassidinae,
feeding niches, 224
Hispinae, 224
trophic selection, 224
Zingiberales, 224
caterpillar,
oxidative stress, 1113
phenolic compounds, 1113
superoxide dismutase, 1113
tannin, 1113
cellulose,
development, 45
pH, 45
stable fly, 45
survival, 45
Centaurea stoebe,
Larinus spp., 510
seed bank, 510
seed production, 510
Urophora affinis, 510
Chamaemyiidae,
Adelges tsugae, 1477
parasitoids, 1477
predators, 1477
Tsuga heterophylla, 1477
chemical defense,
alfalfa, 947
arthropods, 947
diversity cascades, 947
saponins, 947
cherry fruit,
cherry leaves, 1086
nitrogen, 1086
Rhagoletis indifferens, 1086
sugar, 1086
cherry leaves,
cherry fruit, 1086
nitrogen, 1086
Rhagoletis indifferens, 1086
sugar, 1086
Chiapas,
intraguild predation, 181
Mexico, 181
multipredator effect, 181
resource partitioning, 181
trophic interaction, 181
Chironomidae,
secondary production, 1162
sex dimorphism, 1162
sex ratios, 1162
tropical stream, 1162
chitinase,
ecological interactions, 1582
herbivory, 1582
silver birch, 1582
transgenic tree, 1582
choice tests,
glassy-winged sharpshooter, 787
regulated water deficit, 787
Xylella fastidiosa, 787
Choristoneura rosaceana, 366, 679
Cydia pomonella, 679
delayed mating, 679
field trapping, 366
flight tunnel, 366
mating disruption, 679
obliquebanded leafroller, 366
pheromone compounds, 366
population dynamics, 679
Chrysopa pallens,
Aphis gossypii, 263
transgenic *Bt* cotton, 263
tri-tropic interaction, 263
Chrysoperla externa,
cypermethrin, 1252
endosulfan, 1252
methoxyfenozide, 1252
spinosad, 1252
Cirrospilus neotropicus,
Ageniaspis citricola, 1025
citrus leafminer, 1025
host selection, 1025
tritrophic interactions, 1025
citrus leafminer,
Ageniaspis citricola, 1025
Cirrospilus neotropicus, 1025
host selection, 1025
tritrophic interactions, 1025
classical biological control,
Adelges tsugae, 1498
exclusion cage, 1498
field insectary, 1498
synchrony, 1498
classification and regression tree analysis,
climate, 281
disturbance ecology, 281
Engelmann spruce, 281
spruce beetle, 281
climate,
classification and regression tree analysis, 281
disturbance ecology, 281
Engelmann spruce, 281
spruce beetle, 281
climate change,
Dendroctonus frontalis, 650
outbreaks, 650
Thanasimus dubius, 650
upper lethal temperature, 650
climate matching,
ambrosia beetle, 442
cost-weighted distance, 442
redbay, 442
Xyleborus glabratus, 442
Coccinellidae, 143, 1258
ant attendance, 1258
ant-hemipteran mutualism, 143
coffee, 143
Mexico, 143
Monomorium, 1258
myrmecophily, 143
soybean aphid, 1258
coevolution,
Agrilus planipennis, 242
host plant resistance, 242
invasive species, 242
wood borers, 242

coffee,
ant-hemipteran mutualism, 143
Coccinellidae, 143
Mexico, 143
myrmecophily, 143
cold,
migration behavior, 1389
Mythimna separata, 1389
sensitive stage, 1389
starvation, 1389
cole crops,
seasonal development, 907
swede midge, 907
Coleomegilla maculata, 315, 323, 525
Cry1Ab endotoxin, 525
dietary mixing, 525
food deprivation, 315, 323
Harmonia axyridis, 315, 323, 525
Hippodamia convergens, 315, 323
life history traits, 315, 323
zoophytophagy, 525
collective movement,
Anabrus simplex, 889
hopper band, 889
katydid, 889
radiotransmitter, 889
colonization,
aphid-pathogenic fungi, 1264
dissemination, 1264
host dispersal flight, 1264
parasitoids, 1264
common cutworm,
Bt cotton, 1043
population dynamics, 1043
survival, 1043
community structure,
conservation, 1224
moth, 1224
spatial structure, 1224
timber harvest, 1224
compensation,
Diatraea saccharalis, 796
injury, 796
rice, 796
sugarcane borer, 796
compensatory photosynthesis,
defoliation, 1332
folivory, 1332
gas exchange, 1332
plant–insect interaction, 1332
competition, 741, 1307
Argentine ant, 741
bagworm, 1307
host discrimination, 1307
host feeding, 1307
Linepithema humile, 741
odorous house ant, 741
parasitoid, 1307
Tapinoma sessile, 741
competitive attraction,
apple, 1136
Cydia pomonella, 1136
mating disruption, 1136
pest management, 1136
completeness of inventories,
beta diversity, 1488
Calathus, 1488
Iberian Peninsula, 1488
species richness, 1488
connectivity,
dispersal, 897
Lepidoptera, 897
matrix habitat, 897
metapopulation, 897
Conotrachelus nenuphar,
interstrain matings, 1208
reproductive incompatibility, 1208
conservation,
community structure, 1224
moth, 1224
spatial structure, 1224
timber harvest, 1224
conservation biological control,
beetle banks, 478
habitat manipulation, 478
insectary plants, 478
native plants, 478
contiguous habitats,
Argentina, 419
Humid Chaco ecoregion, 419
Mburucuya National Park, 419
spider diversity, 419
conventional versus sustainable agronomic practices,
epigeic predators, 1214
ground beetles, 1214
cooling rate,
body size, 1074, 1074
cooling rate interaction, 1074
fire ant, 1074
supercooling point, 1074
cooling rate interaction,
body size, 1074, 1074
cooling rate, 1074
fire ant, 1074
supercooling point, 1074
α-copaene,
calamenene, 1403
exotic species, 1403
Cormack-Jolly-Seber,
capture-recapture, 1145
survival, 1145
vagility, 1145
corridor,
Carabidae, 725
edge effects, 725
fragmentation, 725
ground beetles, 725
cost-weighted distance,
ambrosia beetle, 442
climate matching, 442
redbay, 442
Xyleborus glabratus, 442
costs of resistance,
Drosophila, 1099
ectoparasitism, 1099
embryo mortality, 1099
Macrocheles mites, 1099
Cotinis nitida L.,
grapes, 608
green June beetle, 608
Japanese beetle, 608
Popillia japonica Newman, 608
cotton, 1396, 1410
Anthonomus grandis grandis, 1396
attractant, 1410
boll weevil, 1396
natural enemy, 1410
photoperiod, 1396
plant volatile, 1410
cotton cultivar,
Aphis gossypii, 29
elevated CO_2, 29
Propylaea japonica, 29
tri-trophic interaction, 29
cotton gin trash,
behavior, 1373
seasonal dynamics, 1373
soil moisture preference, 1373
cottonseed,
Gossypium hirsutum, 1081
(+)-gossypol, 1081
tobacco budworm, 1081
courtship,
parasitoid wasp, 782
receptivity, 782
signaling, 782
vestigial, 782
crop rotation,
cabbage root fly, 409
Delia radicum, 409
spatial dynamics, 409
spatial ecology, 409
crude fat,
insecticide, 206
Nilaparvata lugens, 206
reproductive rate, 206
soluble sugar, 206
Cry1Ab endotoxin,
Coleomegilla maculata, 525
dietary mixing, 525
Harmonia axyridis, 525
zoophytophagy, 525
Cry1Ac level,
population dynamics, 1361
Sesamia inferens, 1361
transgenic rice lines, 1361

Cry1F protein,
Bt accumulation in soil, 255
environmental risk, 255
enzyme-linked immunosorbent assay, 255
transgenic corn, 255
Cry3Bb1,
biotechnology, 838
nontarget organisms, 838
potassium arsenate, 838
protease inhibitor, 838
cryobiology,
aerobic metabolism, 1069
freeze tolerance, 1069
glycerol, 1069
Hypercompe scribonia, 1069
Curculio caryae,
Beauveria bassiana, 162
biological control, 162
pecan, 162
Curculionidae,
Bostrichidae, 1300
host finding, 1300
learning, 1300
Pteromalidae, 1300
Cydia pomonella, 679, 1136
apple, 1136
Choristoneura rosaceana, 679
competitive attraction, 1136
delayed mating, 679
mating disruption, 679, 1136
pest management, 1136
population dynamics, 679
Cynodon dactylon,
Antonina graminis, 487
honeydew, 487
invasive species, 487
Solenopsis invicta, 487
cypermethrin,
Chrysoperla externa, 1252
endosulfan, 1252
methoxyfenozide, 1252
spinosad, 1252

D

D. frontalis,
amplified fragment length polymorphism, 817
Dendroctonus brevicomis, 817
genetic differentiation, 817
Ips pini, 817
Dacini,
adult life history, 307
host-plant effect, 307
preimaginal development, 307
temperature, 307
daily variations,
animal foraging, 857
harvester ants, 857
soil temperature, 857
trail system, 857
dairy manure,
black soldier fly larvae, 1525
Escherichia coli, 1525
Hermetia illucens, 1525
Stratiomyidae, 1525
day-flying moths,
butterflies, 1313
estimating species richness, 1313
sample completeness, 1313
sampling fraction, 1313
defoliation,
compensatory photosynthesis, 1332
folivory, 1332
gas exchange, 1332
plant–insect interaction, 1332
degree day, 57, 568
Dendroctonus, 57
Ips, 57
larval development, 568
life table, 568
management, 568
Pinus ponderosa, 57
temperature, 57
relative humidity, 568
delayed mating,
Choristoneura rosaceana, 679
Cydia pomonella, 679
mating disruption, 679
population dynamics, 679
Delia platura,
Brassica napus, 1451
Brassica rapa, 1451
Delia radicum, 1451
logistic model, 1451
Delia radicum, 409, 1451
Brassica napus, 1451
Brassica rapa, 1451
cabbage root fly, 409
crop rotation, 409
Delia platura, 1451
logistic model, 1451
spatial dynamics, 409
spatial ecology, 409
Dendroctonus,
degree day, 57
Ips, 57
Pinus ponderosa, 57
temperature, 57
Dendroctonus,
abundance, 94
flight seasonality, 94
Ips, 94
Pinus ponderosa, 94
Dendroctonus brevicomis,
amplified fragment length polymorphism, 817
D. frontalis, 817
genetic differentiation, 817
Ips pini, 817
Dendroctonus frontalis, 271, 650
climate change, 650
insect pest, 271
microsatellites, 271
outbreaks, 650
population structure, 271
southern pine beetle, 271
Thanasimus dubius, 650
upper lethal temperature, 650
Dendroctonus ponderosae,
4-allylanisole, 871
estragole, 871
mountain pine beetle, 871
Pinus ponderosa, 871
Dendroctonus pseudotsugae,
Douglas-fir beetle, 850
fatreserves, 850
insect flight, 850
Soxhlet extraction, 850
development, 45, 340, 630, 660, 1105
acalypha, 630
cellulose, 45
hibiscus, 630
life table, 340, 1105
Liposcelis badia, 1105
parthenium, 630
pH, 45
phenology simulation model, 660
Phytoseiidae, 340
plumeria, 630
pollen, 340
Scotinophara lurida, 660
stable fly, 45
stored product, 1105
survival, 45
temperature, 660, 1105
Typhlodromus foenilis, 340
developmental rate, 38, 348, 862, 1381
ArcGIS mapping, 862
biological control, 862
Brassica oleracea, 38
Diadegma anurum, 38
embryo development, 1381
Episimus utilis, 862
invasive species, 348
modeling, 1381
nonlinear model, 348
Plutella xylostella, 38
Schinus terebinthifolius, 862
Sitona discoideus, 1381
stink bug, 348
temperature, 38, 1381
developmental time,
Helicoverpa armigera, 16
life table, 16
survival, 16
temperature, 16

Diabrotica,
DNA markers, 293
microsatellites, 293
population genetics, 293
western corn rootworm, 293
Diabrotica barberi,
alternate hosts, 247
prairie grasses, 247
resistance management, 247
Diabrotica virgifera virgifera,
phenology, 1558
root toughness, 1558
Diadegma anurum,
Brassica oleracea, 38
development rate, 38
Plutella xylostella, 38
temperature, 38
Dialeurodes citri,
beneficial arthropods, 555
Lepidosaphes beckii, 555
Panonychus citri, 555
Satsuma mandarin, 555
diapause,
apple, 1154
Rhagoletis pomonella, 1154
Diaprepes root weevil,
egg parasitoid, 1565
host plants, 1565
searching speed, 1565
trichomes, 1565
Diatraea saccharalis,
compensation, 796
injury, 796
rice, 796
sugarcane borer, 796
diet switching,
beneficial acclimation hypothesis, 1003
calorimetry, 1003
dietary acclimation hypothesis, 1003
nutritional ecology, 1003
dietary acclimation hypothesis,
beneficial acclimation hypothesis, 1003
calorimetry, 1003
diet switching, 1003
nutritional ecology, 1003
dietary mixing,
Coleomegilla maculata, 525
Cry1Ab endotoxin, 525
Harmonia axyridis, 525
zoophytophagy, 525
dietary phosphorus limitation to grasshoppers,
ecological stoichiometry, 333
multiple nutrients, 333
nutritional ecology, 333
nutritional indices, 333
Diptera, 505, 876
Anastrepha obliqua, 876
biological management, 505
calling behavior, 876
entomopathogenic bacterium, 505
male volatiles, 876
microbial insecticide, 505
Tephritidae, 876
treatments, 505
dipterocarp forest,
α diversity, 1461
β diversity, 1461
effective number of species, 1461
ethanol trap, 1461
flight intercept trap, 1461
dispersal, 301, 636, 897, 1280, 1291
Aphis glycines, 301
Cactoblastis cactorum, 1291
connectivity, 897
environmental condition, 301
female flight capability, 636
flight, 1291
flight ability, 301
flight behavior, 1280
flight mill, 301, 1280
genetic variation, 636
gypsy moth, 636
Lepidoptera, 897
Lymantria dispar, 636
matrix habitat, 897
metapopulation, 897
migration, 1280
Ostrinia nubilalis, 1280
overwintering, 1291
seasonality, 1291
dissemination,
aphid-pathogenic fungi, 1264
colonization, 1264
host dispersal flight, 1264
parasitoids, 1264
distribution patterns,
canopy lichens, 464
microhabitat specificity, 464
montane forests, 464
Oribatid mites, 464
disturbance ecology,
classification and regression tree analysis, 281
climate, 281
Engelmann spruce, 281
spruce beetle, 281
diversity cascades,
alfalfa, 947
arthropods, 947
chemical defense, 947
saponins, 947
DNA markers,
Diabrotica, 293
microsatellites, 293
population genetics, 293
western corn rootworm, 293
Douglas-fir beetle,
Dendroctonus pseudotsugae, 850
fatreserves, 850
insect flight, 850
Soxhlet extraction, 850
Drosophila,
costs of resistance, 1099
ectoparasitism, 1099
embryo mortality, 1099
Macrocheles mites, 1099
dusky wireworm,
Agriotes obscurus, 534
insecticide, 534
repellency, 534
dynamic head space sampling,
electroantennogram, 1416
sex pheromone, 1416
tea mosquito, 1416
wind tunnel, 1416

E

ecological assembly,
Black Belt Prairie, 453
Formicidae, 453
Mississippi, 453
Solenopsis, 453
ecological interactions,
chitinase, 1582
herbivory, 1582
silver birch, 1582
transgenic tree, 1582
ecological stoichiometry,
dietary phosphorus limitation to grasshoppers, 333
multiple nutrients, 333
nutritional ecology, 333
nutritional indices, 333
ecosystem engineering,
habitat fragmentation, 938
sika deer, 938
vegetation structure, 938
ectoparasitism,
costs of resistance, 1099
Drosophila, 1099
embryo mortality, 1099
Macrocheles mites, 1099
edge effects,
Carabidae, 725
corridor, 725
fragmentation, 725
ground beetles, 725
effective number of species,
ga diversity, 1461
gb diversity, 1461
dipterocarp forest, 1461
ethanol trap, 1461
flight intercept trap, 1461
egg maturation,
invasive species, 1200
Pierce's disease, 1200
Xylella fastidiosa, 1200

egg parasitoid,
Diaprepes root weevil, 1565
host plants, 1565
searching speed, 1565
trichomes, 1565
electrical penetration graph,
Frankliniella fusca, 213
Frankliniella occidentalis, 213
tomato, 213
electroantennogram,
dynamic head space sampling, 1416
sex pheromone, 1416
tea mosquito, 1416
wind tunnel, 1416
electroantennography,
cadaverine, 1119
Caribbean fruit fly, 1119
putrescine, 1119
elevated carbon dioxide,
fecundity, 601
Glycine max, 601
longevity, 601
Popillia japonica, 601
elevated CO_2,
Aphis gossypii, 29
cotton cultivar, 29
Propylaea japonica, 29
tri-trophic interaction, 29
embryo development,
developmental rate, 1381
modeling, 1381
Sitona discoideus, 1381
temperature, 1381
embryo mortality,
costs of resistance, 1099
Drosophila, 1099
ectoparasitism, 1099
Macrocheles mites, 1099
emerald ash borer,
forest insect, 1344
insect gut, 1344
invasive species, 1344
symbiont, 1344
encapsulation,
body size, 51
habitat quality, 51
immunity, 51
trade-off, 51
Encarsia spp.,
Bemisia tabaci biotype B, 391
biological control, 391
Eretmocerus spp., 391
sweetpotato whitefly, 391
endosulfan,
Chrysoperla externa, 1252
cypermethrin, 1252
methoxyfenozide, 1252
spinosad, 1252
Engelmann spruce,
classification and regression tree analysis, 281
climate, 281
disturbance ecology, 281
spruce beetle, 281
entomopathogenic bacterium,
biological management, 505
Diptera, 505
microbial insecticide, 505
treatments, 505
entomopathogenic nematodes,
pupation, 845
soil moisture, 845
soil texture, 845
environmental condition,
Aphis glycines, 301
dispersal, 301
flight ability, 301
flight mill, 301
environmental risk,
Bt accumulation in soil, 255
Cry1F protein, 255
enzyme-linked immunosorbent assay, 255
transgenic corn, 255
enzyme-linked immunosorbent assay, 255, 990
Bt accumulation in soil, 255
Cry1F protein, 255
environmental risk, 255
gut content analysis, 990
monoclonal antibody, 990
predator–prey interactions, 990
sampling protocols, 990
transgenic corn, 255
7-*epi*-sesquithujene,
bark volatiles, 356
Buprestidae, 356
Manuka oil, 356
Phoebe oil, 356
epidemiology,
insect dispersal, 79
tomato spotted wilt virus, 79
epigeic predators,
conventional versus sustainable agronomic practices, 1214
ground beetles, 1214
Episimus utilis,
ArcGIS mapping, 862
biological control, 862
developmental rate, 862
Schinus terebinthifolius, 862
Eretmocerus spp.,
Bemisia tabaci biotype B, 391
biological control, 391
Encarsia spp., 391
sweetpotato whitefly, 391
Escherichia coli,
black soldier fly larvae, 1525
dairy manure, 1525
Hermetia illucens, 1525
Stratiomyidae, 1525
estimating species richness,
butterflies, 1313
day-flying moths, 1313
sample completeness, 1313
sampling fraction, 1313
estragole,
4-allylanisole, 871
Dendroctonus ponderosae, 871
mountain pine beetle, 871
Pinus ponderosa, 871
ethanol trap,
α diversity, 1461
β diversity, 1461
dipterocarp forest, 1461
effective number of species, 1461
flight intercept trap, 1461
Euproctis chrysorrhoea,
forest defoliator, 1429
invasive species, 1429
overwintering mortality, 1429
population collapse, 1429
European corn borer,
genetically engineered corn, 615
insect resistance management, 615
maize, 615
Euschistus conspersus,
mate choice, 192
mating behavior, 192
parental investment, 192
reproductive behavior, 192
exclusion cage,
Adelges tsugae, 1498
classical biological control, 1498
field insectary, 1498
synchrony, 1498
exo-brevicomin,
interspecific communication, 70
pheromones, 70
reproductive isolation, 70
trapping, 70
exotic species,
calamenene, 1403
α-copaene, 1403
experimental design,
Bacillus thuringiensis, 1
genetically modified, 1
risk assessment, 1
extrafloral,
anthrone, 996
nectar, 996
nutrition, 996
spider, 996

F

fatreserves,
Dendroctonus pseudotsugae, 850
Douglas-fir beetle, 850
insect flight, 850
Soxhlet extraction, 850
fecundity, 520, 586, 601
elevated carbon dioxide, 601
Glycine max, 601
insect/plant interactions, 520
instrinsic rate of natural increase, 586
longevity, 586, 601
net reproductive rate, 586
oviposition, 520
Popillia japonica, 601
feeding behavior,
feeding ecology, 172
protein, 172
Rhagoletis indifferens, 172
sugar, 172
feeding choice tests,
food mixing, 1323
host plant selection, 1323
leaffooted bugs, 1323
passion vines, 1323
feeding ecology,
feeding behavior, 172
protein, 172
Rhagoletis indifferens, 172
sugar, 172
feeding niches,
Cassidinae, 224
Hispinae, 224
trophic selection, 224
Zingiberales, 224
female flight capability,
dispersal, 636
genetic variation, 636
gypsy moth, 636
Lymantria dispar, 636
field insectary,
Adelges tsugae, 1498
classical biological control, 1498
exclusion cage, 1498
synchrony, 1498
field trapping,
Choristoneura rosaceana, 366
flight tunnel, 366
obliquebanded leafroller, 366
pheromone compounds, 366
finite rate of increase,
generation time, 1538
net reproductive rate, 1538
plant host abundance, 1538
tobacco budworm, 1538
fire ant,
body size, 1074, 1074
cooling rate, 1074
cooling rate interaction, 1074
supercooling point, 1074
flight,
Cactoblastis cactorum, 1291
dispersal, 1291
overwintering, 1291
seasonality, 1291
flight ability,
Aphis glycines, 301
dispersal, 301
environmental condition, 301
flight mill, 301
flight behavior, 774, 1280
autumn migration, 774
dispersal, 1280
flight mill, 1280
light trapping, 774
migration, 1280
Mythimna separata, 774
Ostrinia nubilalis, 1280
radar observation, 774
flight capacity,
mating status, 374
ovarian development, 374
flight intercept trap,
α diversity, 1461
β diversity, 1461
dipterocarp forest, 1461
effective number of species, 1461
ethanol trap, 1461
flight mill, 301, 1280
Aphis glycines, 301
dispersal, 301, 1280
environmental condition, 301
flight ability, 301
flight behavior, 1280
migration, 1280
Ostrinia nubilalis, 1280
flight seasonality,
abundance, 94
Dendroctonus, 94
Ips, 94
Pinus ponderosa, 94
flight tunnel,
Choristoneura rosaceana, 366
field trapping, 366
obliquebanded leafroller, 366
pheromone compounds, 366
folivory,
compensatory photosynthesis, 1332
defoliation, 1332
gas exchange, 1332
plant–insect interaction, 1332
food deprivation, 315, 323
Coleomegilla maculata, *315*, *323*
Harmonia axyridis, 315, 323
Hippodamia convergens, 315, 323
life history traits, 315, 323
food limitation,
grasshoppers, 686
insect population dynamics, 686
precipitation, 686
weather conditions, 686
food mixing,
feeding choice tests, 1323
host plant selection, 1323
leaffooted bugs, 1323
passion vines, 1323
forest defoliator,
Euproctis chrysorrhoea, 1429
invasive species, 1429
overwintering mortality, 1429
population collapse, 1429
forest insect, 1174, 1344
emerald ash borer, 1344
habitat structure, 1174
insect gut, 1344
invasive species, 1174, 1344
Lymantria dispar, 1174
natural enemies, 1174
symbiont, 1344
Formicidae,
Black Belt Prairie, 453
ecological assembly, 453
Mississippi, 453
Solenopsis, 453
fragmentation,
Carabidae, 725
corridor, 725
edge effects, 725
ground beetles, 725
Frankliniella fusca,
electrical penetration graph, 213
Frankliniella occidentalis, 213
tomato, 213
Frankliniella occidentalis,
electrical penetration graph, 213
Frankliniella fusca, *213*
tomato, 213
Fraxinus,
host range, 230
host selection, 230
invasive pest, 230
oviposition preference, 230
freeze tolerance,
aerobic metabolism, 1069
cryobiology, 1069
glycerol, 1069
Hypercompe scribonia, 1069
fungal pathogen,
fungal repellence, 755
integrated pest management, 755
landscape mulch management, 755
subterranean termites, 755

fungal repellence,
fungal pathogen, 755
integrated pest management, 755
landscape mulch management, 755
subterranean termites, 755

G

gas exchange,
compensatory photosynthesis, 1332
defoliation, 1332
folivory, 1332
plant–insect interaction, 1332
gene flow,
Bacillus thuringiensis, 1354
Mediterranean corn borer, 1354
RAPD, 1354
transgenic maize, 1354
generalist predators,
agro-ecology, 121
beetle communities, 121
biological control, 121
natural enemies, 121
generation time,
finite rate of increase, 1538
net reproductive rate, 1538
plant host abundance, 1538
tobacco budworm, 1538
genetic differentiation,
amplified fragment length polymorphism, 817
D. frontalis, 817
Dendroctonus brevicomis, 817
Ips pini, 817
genetic variation,
dispersal, 636
female flight capability, 636
gypsy moth, 636
Lymantria dispar, 636
genetically engineered corn,
European corn borer, 615
insect resistance management, 615
maize, 615
genetically modified,
Bacillus thuringiensis, 1
experimental design, 1
risk assessment, 1
geographical distribution,
Aphidius ervi, 472
phenology, 472
population dynamics, 472
Praon unicum, 472
geostatistics,
Lobesia botrana, 382
olive grove, 382
vineyard, 382
glassy-winged sharpshooter, 787, 925, 1049
Auchenorrhyncha, 925
behavior, 1049
choice tests, 787
Hexapoda, 925
leafhopper, 925
management, 1049
Pierce's disease, 925
regulated water deficit, 787
Xylella fastidiosa, 787, 1049
global warming,
phenology, 696
Pieris rapae, 696
precipitation, 696
temperature, 696
glycerol,
aerobic metabolism, 1069
cryobiology, 1069
freeze tolerance, 1069
Hypercompe scribonia, 1069
Glycine max,
elevated carbon dioxide, 601
fecundity, 601
longevity, 601
Popillia japonica, 601
Gossypium hirsutum,
cottonseed, 1081
(+)-gossypol, 1081
tobacco budworm, 1081
(+)-gossypol,
cottonseed, 1081
Gossypium hirsutum, 1081
tobacco budworm, 1081
grapes,
Cotinis nitida L, 608
green June beetle, 608
Japanese beetle, 608
Popillia japonica Newman, 608
grasshoppers,
food limitation, 686
insect population dynamics, 686
precipitation, 686
weather conditions, 686
green June beetle,
Cotinis nitida L, 608
grapes, 608
Japanese beetle, 608
Popillia japonica Newman, 608
ground beetles, 725, 1214
Carabidae, 725
conventional versus sustainable agronomic practices, 1214
corridor, 725
edge effects, 725
epigeic predators, 1214
fragmentation, 725
growth rate,
Alatae, 1573
host choice, 1573
plant pathogen, 1573
virus vector, 1573
gut content analysis,
enzyme-linked immunosorbent assay, 990
monoclonal antibody, 990
predator–prey interactions, 990
sampling protocols, 990
gypsy moth,
dispersal, 636
female flight capability, 636
genetic variation, 636
Lymantria dispar, 636

H

habitat fragmentation,
ecosystem engineering, 938
sika deer, 938
vegetation structure, 938
habitat management, 707, 719
bumble bee, 707, 719
pollination, 707, 719
prairie plants, 707, 719
restoration, 707, 719
habitat manipulation,
beetle banks, 478
conservation biological control, 478
insectary plants, 478
native plants, 478
habitat quality,
body size, 51
encapsulation, 51
immunity, 51
trade-off, 51
habitat structure,
forest insects, 1174
invasive species, 1174
Lymantria dispar, 1174
natural enemies, 1174
haplodiploidy,
arrhenotoky, 1422
thelytoky, 1422
Thysanoptera, 1422
Wolbachia, 1422
Harmonia axyridis, 315, 323, 525, 973
Aphis glycines, 973
buckthorn, 973
Coleomegilla maculata, 315, 323
Coleomegilla maculata, 525
Cry1Ab endotoxin, 525
dietary mixing, 525
food deprivation, 315, 323
Hippodamia convergens, 315, 323
life history traits, 315, 323
preference assay, 973
soybean aphid, 973

zoophytophagy, 525
harvester ants,
animal foraging, 857
daily variations, 857
soil temperature, 857
trail system, 857
Helianthus annuus,
arbuscular mycorrhizal fungi, 1017
sunflower beetle, 1017
trait-mediated indirect effects, 1017
Zygogramma exclamationis, 1017
Helicoverpa armigera,
developmental time, 16
life table, 16
survival, 16
temperature, 16
herbivory,
chitinase, 1582
ecological interactions, 1582
silver birch, 1582
transgenic tree, 1582
Hermetia illucens,
black soldier fly larvae, 1525
dairy manure, 1525
Escherichia coli, 1525
Stratiomyidae, 1525
Heterorhabditis bacteriophora,
reproduction, 1272
Steinernema feltiae, 1272
thermal adaptation, 1272
virulence, 1272
Hexapoda,
Auchenorrhyncha, 925
glassy-winged sharpshooter, 925
leafhopper, 925
Pierce's disease, 925
hibiscus,
acalypha, 630
development, 630
parthenium, 630
plumeria, 630
Hippodamia convergens, 315, 323
Coleomegilla maculata, 315, 323
food deprivation, 315, 323
Harmonia axyridis, 315, 323
life history traits, 315, 323
Hispinae,
Cassidinae, 224
feeding niches, 224
trophic selection, 224
Zingiberales, 224
Histiogaster,
bark beetles, 956
Hyphomycetes, 956
Leptographium, 956
symbiosis, 956
honey bee, 546, 808
Apis mellifera, 808
incidents, 546
invasive species, 808
mortality, 546
native plant, 808
pesticides, 546
risk assessment, 546
Triteleia laxa, 808
honeydew,
Antonina graminis, 487
Cynodon dactylon, 487
invasive species, 487
Solenopsis invicta, 487
hopper band,
Anabrus simplex, 889
collective movement, 889
katydid, 889
radiotransmitter, 889
host choice,
Alatae, 1573
growth rate, 1573
plant pathogen, 1573
virus vector, 1573
host discrimination,
bagworm, 1307
competition, 1307
host feeding, 1307
parasitoid, 1307
host dispersal flight,
aphid-pathogenic fungi, 1264
colonization, 1264
dissemination, 1264
parasitoids, 1264
host feeding,
bagworm, 1307
competition, 1307
host discrimination, 1307
parasitoid, 1307
host finding,
Bostrichidae, 1300
Curculionidae, 1300
learning, 1300
Pteromalidae, 1300
host plant,
biological control, 1185
Diaprepes root weevil, 1565
egg parasitoid, 1565
life history, 1185
Lixinae, 1185
phenology, 1185
searching speed, 1565
trichomes, 1565
host plant resistance,
Agrilus planipennis, 242
coevolution, 242
invasive species, 242
wood borers, 242
host plant selection, 1323, 1548
feeding choice tests, 1323
food mixing, 1323
Japanese beetle, 1548
leaffooted bugs, 1323
passion vines, 1323
plant defense theory, 1548
polyphagy, 1548
Popillia japonica, 1548
host range, 230, 1231
Fraxinus, 230
host selection, 230
host-instar utilization, 1231
hyperparasitism, 1231
idiobiont/koinobiont mode, 1231
invasive pest, 230
oviposition preference, 230
parasitism, 1231
host selection, 230, 1025
Ageniaspis citricola, 1025
Cirrospilus neotropicus, 1025
citrus leafminer, 1025
Fraxinus, 230
host range, 230
invasive pest, 230
oviposition preference, 230
tritrophic interactions, 1025
host suitability,
Anagrus nilaparvatae, 748
infestation level, 748
Nilaparvata lugens, 748
tritrophic interaction, 748
host-instar utilization,
host range, 1231
hyperparasitism, 1231
idiobiont/koinobiont mode, 1231
parasitism, 1231
host–plant effect,
adult life history, 307
Dacini, 307
host–plant effect, 307
preimaginal development, 307
temperature, 307
huanglongbing,
phenology, 914
sampling, 914
sticky traps, 914
trapping, 914
Humid Chaco ecoregion,
Argentina, 419
contiguous habitats, 419
Mburucuya National Park, 419
spider diversity, 419
Hypercompe scribonia,
aerobic metabolism, 1069
cryobiology, 1069
freeze tolerance, 1069
glycerol, 1069
hyperparasitism,
host range, 1231
host-instar utilization, 1231
idiobiont/koinobiont mode, 1231
parasitism, 1231

Hyphomycetes,
bark beetles, 956
Histiogaster, 956
Leptographium, 956
symbiosis, 956

I

Iberian Peninsula,
beta diversity, 1488
Calathus, 1488
completeness of inventories, 1488
species richness, 1488
idiobiont/koinobiont mode,
host range, 1231
host-instar utilization, 1231
hyperparasitism, 1231
parasitism, 1231
immunity,
body size, 51
encapsulation, 51
habitat quality, 51
trade-off, 51
inbreeding,
inclusive fitness, 883
neotenic reproductives, 883
reproductive plasticity, 883
reproductive skew, 883
incidents,
honey bees, 546
mortality, 546
pesticides, 546
risk assessment, 546
inclusive fitness,
inbreeding, 883
neotenic reproductives, 883
reproductive plasticity, 883
reproductive skew, 883
indirect competition,
biological control, 734
mutualism, 734
scale insects, 734
shade coffee, 734
infestation level,
Anagrus nilaparvatae, 748
host suitability, 748
Nilaparvata lugens, 748
tritrophic interaction, 748
injury,
compensation, 796
Diatraea saccharalis, 796
rice, 796
sugarcane borer, 796
insect,
integrated pest management, 575
nitrogen, 575
yield, 575
insect dispersal,
epidemiology, 79
tomato spotted wilt virus, 79
insect flight,
Dendroctonus pseudotsugae, 850
Douglas-fir beetle, 850
fatreserves, 850
Soxhlet extraction, 850
insect gut,
emerald ash borer, 1344
forest insect, 1344
invasive species, 1344
symbiont, 1344
insect pest,
Dendroctonus frontalis, 271
microsatellites, 271
population structure, 271
southern pine beetle, 271
insect/plant interactions,
fecundity, 520
oviposition, 520
insect population dynamics,
food limitation, 686
grasshoppers, 686
precipitation, 686
weather conditions, 686
insect resistance management, 615, 1514
Bemisia tabaci, 1514
European corn borer, 615
genetically engineered corn, 615
maize, 615
modeling, 1514
operational and environmental factors, 1514
pyriproxyfen, 1514
insect vectors,
alternate weed hosts, 592
vector performance, 592
vector reservoir, 592
viral inoculum source, 592
insectary plants,
beetle banks, 478
conservation biological control, 478
habitat manipulation, 478
native plants, 478
insecticide, 206, 534
Agriotes obscurus, 534
crude fat, 206
dusky wireworm, 534
Nilaparvata lugens, 206
repellency, 534
reproductive rate, 206
soluble sugar, 206
insidious flower bug,
preference, 964
prey consumption, 964
soybean aphid, 964
soybean thrips, 964
insolation,
nest orientation, 198
remote sensing, 198
thermoregulation, 198
instrinsic rate of natural increase,
fecundity, 586
longevity, 586
net reproductive rate, 586
integrated pest management, 575, 755
fungal pathogen, 755
fungal repellence, 755
insect, 575
landscape mulch management, 755
nitrogen, 575
subterranean termites, 755
yield, 575
interspecific communication,
exo-brevicomin, 70
pheromones, 70
reproductive isolation, 70
trapping, 70
interstrain matings,
Conotrachelus nenuphar, 1208
reproductive incompatibility, 1208
intraguild predation,
Chiapas, 181
Mexico, 181
multipredator effect, 181
resource partitioning, 181
trophic interaction, 181
invasive pest,
Fraxinus, 230
host range, 230
host selection, 230
oviposition preference, 230
invasive species, 242, 348, 487, 808, 1174, 1200, 1344, 1429
Agrilus planipennis, 242
Antonina graminis, 487
Apis mellifera, 808
coevolution, 242
Cynodon dactylon, 487
development rate, 348
egg maturation, 1200
emerald ash borer, 1344
Euproctis chrysorrhoea, 1429
forest defoliator, 1429
forest insect, 1174, 1344
habitat structure, 1174
honey bee, 808
honeydew, 487
host plant resistance, 242
insect gut, 1344
Lymantria dispar, 1174
native plant, 808
natural enemies, 1174
nonlinear model, 348
overwintering mortality, 1429
Pierce's disease, 1200
population collapse, 1429
Solenopsis invicta, 487
stink bug, 348

symbiont, 1344
Triteleia laxa, 808
wood borers, 242
Xylella fastidiosa, 1200
Ips, 57, 94
abundance, 94
degree day, 57
Dendroctonus, 57, 94
flight seasonality, 94
Pinus ponderosa, 57, 94
temperature, 57
Ips pini, 150, 817
amplified fragment length polymorphism, 817
D. frontalis, 817
Dendroctonus brevicomis, 817
genetic differentiation, 817
microbe interactions, 150
parasitoid behavior, 150
predator behavior, 150
semiochemicals, 150
Isoptera,
breeding system, 400
microsatellite genotyping, 400
morphometry, 400
Rhinotermitidae, 400

J

Japanese beetle, 608, 1548
Cotinis nitida L, 608
grapes, 608
green June beetle, 608
host plant selection, 1548
plant defense theory, 1548
polyphagy, 1548
Popillia japonica, 1548
Popillia japonica Newman, 608

K

katydid,
Anabrus simplex, 889
collective movement, 889
hopper band, 889
radiotransmitter, 889

L

landscape mulch management,
fungal pathogen, 755
fungal repellence, 755
integrated pest management, 755
subterranean termites, 755
Laricobius nigrinus,
Adelges tsugae, 494
Leucopis argenticollis, 494
Leucopis atrifacies, 494
Tsuga heterophylla, 494
Larinus spp.,
Centaurea stoebe, 510
seed bank, 510
seed production, 510
Urophora affinis, 510
larval development,
degree-days, 568
life table, 568
management, 568
relative humidity, 568
larval parasitoid,
Bactrocera oleae (Gmelin), 764
Olea europaea L., 764
leaffooted bugs,
feeding choice tests, 1323
food mixing, 1323
host plant selection, 1323
passion vines, 1323
leafhopper,
ant, 1471
Auchenorrhyncha, 925
glassy-winged sharpshooter, 925
Hexapoda, 925
Pierce's disease, 925
mutualism, 1471
shade, 1471
shelter, 1471
learning,
Bostrichidae, 1300
Curculionidae, 1300
host finding, 1300
Pteromalidae, 1300
Lepidoptera,
connectivity, 897
dispersal, 897
matrix habitat, 897
metapopulation, 897
Lepidosaphes beckii,
beneficial arthropods, 555
Dialeurodes citri, 555
Panonychus citri, 555
Satsuma mandarin, 555
Leptographium,
bark beetles, 956
Histiogaster, 956
Hyphomycetes, 956
symbiosis, 956
Leucopis argenticollis,
Adelges tsugae, 494
Laricobius nigrinus, 494
Leucopis atrifacies, 494
Tsuga heterophylla, 494
Leucopis atrifacies,
Adelges tsugae, 494
Laricobius nigrinus, 494
Leucopis argenticollis, 494
Tsuga heterophylla, 494
life history,
biological control, 1185
host plant, 1185
Lixinae, 1185
phenology, 1185
life history traits, 315, 323
Coleomegilla maculata, *315*, *323*
food deprivation, 315, 323
Harmonia axyridis, 315, 323
Hippodamia convergens, 315, 323
life table, 16, 340, 568, 1105
degree-days, 568
development, 340, 1105
developmental time, 16
Helicoverpa armigera, 16
larval development, 568
Liposcelis badia, 1105
management, 568
Phytoseiidae, 340
pollen, 340
relative humidity, 568
stored product, 1105
survival, 16
temperature, 16, 1105
Typhlodromus foenilis, 340
light trapping,
autumn migration, 774
flight behavior, 774
Mythimna separata, 774
radar observation, 774
Linepithema humile,
Argentine ant, 741
competition, 741
odorous house ant, 741
Tapinoma sessile, 741
Liposcelis badia,
development, 1105
life table, 1105
stored product, 1105
temperature, 1105
Lixinae,
biological control, 1185
host plant, 1185
life history, 1185
phenology, 1185
Lobesia botrana,
geostatistics, 382
olive grove, 382
vineyard, 382
local indicator of spatial autocorrelation,
aerobiology, 87
biological invasions, 87
Lymantria dispar, 87
quantile regression, 87
logistic model,
Brassica napus, 1451
Brassica rapa, 1451
Delia platura, 1451
Delia radicum, 1451
longevity, 586, 601
elevated carbon dioxide, 601
fecundity, 586, 601
Glycine max, 601
instrinsic rate of natural increase, 586
net reproductive rate, 586
Popillia japonica, 601

low volume,
apple, 980
mating disruption, 980
pear, 980
pest management, 980
Lymantria dispar, 87, 636
aerobiology, 87
biological invasions, 87
dispersal, 636
female flight capability, 636
genetic variation, 636
gypsy moth, 636
local indicator of spatial autocorrelation, 87
quantile regression, 87
Lymantria dispar,
forest insects, 1174
habitat structure, 1174
invasive species, 1174
natural enemies, 1174
Lysiphlebus testaceipes,
Schizaphis graminum, 1063
supercooling point, 1063
winter hardiness, 1063

M

Macrocheles mites,
costs of resistance, 1099
Drosophila, 1099
ectoparasitism, 1099
embryo mortality, 1099
maize,
European corn borer, 615
genetically engineered corn, 615
insect resistance management, 615
male volatiles,
Anastrepha obliqua, 876
calling behavior, 876
Diptera, 876
Tephritidae, 876
management, 568, 1049
behavior, 1049
degree-days, 568
glassy-winged sharpshooter, 1049
larval development, 568
life table, 568
relative humidity, 568
Xylella fastidiosa, 1049
Mangifera indica,
Anastrepha obliqua, 110
Anastrepha suspensa, 110
Utetes anastrephae, 110
Manuka oil,
bark volatiles, 356
Buprestidae, 356
7-*epi*-sesquithujene, 356
Phoebe oil, 356
mate choice,
Euschistus conspersus, 192
mating behavior, 192
parental investment, 192
reproductive behavior, 192
mating behavior,
Euschistus conspersus, 192
mate choice, 192
parental investment, 192
reproductive behavior, 192
mating disruption, 679, 980, 1126, 1136
Anomala orientalis, 1126
apple, 980, 1136
Choristoneura rosaceana, 679
competitive attraction, 1136
Cydia pomonella, 679, 1136
delayed mating, 679
low volume, 980
pear, 980
pest management, 980, 1136
population dynamics, 679
sex pheromone, 1126
turfgrass, 1126
(Z) and (E)-7-tetradecen-2-one formulation, 1126
mating status,
flight capacity, 374
ovarian development, 374
matrix habitat,
connectivity, 897
dispersal, 897
Lepidoptera, 897
metapopulation, 897
Mburucuya National Park,
Argentina, 419
contiguous habitats, 419
Humid Chaco ecoregion, 419
spider diversity, 419
Mediterranean corn borer,
Bacillus thuringiensis, 1354
gene flow, 1354
RAPD, 1354
transgenic maize, 1354
metapopulation,
connectivity, 897
dispersal, 897
Lepidoptera, 897
matrix habitat, 897
methoxyfenozide,
Chrysoperla externa, 1252
cypermethrin, 1252
endosulfan, 1252
spinosad, 1252
Mexico, 143, 181
ant-hemipteran mutualism, 143
Chiapas, 181
Coccinellidae, 143
coffee, 143
intraguild predation, 181
multipredator effect, 181
myrmecophily, 143
resource partitioning, 181
trophic interaction, 181
microbe interactions,
Ips pini, 150
parasitoid behavior, 150
predator behavior, 150
semiochemicals, 150
microbial insecticide,
biological management, 505
Diptera, 505
entomopathogenic bacterium, 505
treatments, 505
microhabitat specificity,
canopy lichens, 464
distribution patterns, 464
montane forests, 464
Oribatid mites, 464
microsatellite genotyping,
breeding system, 400
Isoptera, 400
morphometry, 400
Rhinotermitidae, 400
microsatellites, 271, 293
Dendroctonus frontalis, 271
Diabrotica, 293
DNA markers, 293
insect pest, 271
population genetics, 293
population structure, 271
southern pine beetle, 271
western corn rootworm, 293
migration,
dispersal, 1280
flight behavior, 1280
flight mill, 1280
Ostrinia nubilalis, 1280
migration behavior,
cold, 1389
Mythimna separata, 1389
sensitive stage, 1389
starvation, 1389
Mississippi,
Black Belt Prairie, 453
ecological assembly, 453
Formicidae, 453
Solenopsis, 453
modeling, 1381, 1514
Bemisia tabaci, 1514
developmental rate, 1381
embryo development, 1381
insect resistance management, 1514
operational and environmental factors, 1514
pyriproxyfen, 1514
Sitona discoideus, 1381
temperature, 1381

monoclonal antibody,
enzyme-linked immunosorbent assay, 990
gut content analysis, 990
predator–prey interactions, 990
sampling protocols, 990
Monomorium,
ant attendance, 1258
Coccinellidae, 1258
soybean aphid, 1258
montane forests,
canopy lichens, 464
distribution patterns, 464
microhabitat specificity, 464
Oribatid mites, 464
morphometry,
breeding system, 400
Isoptera, 400
microsatellite genotyping, 400
Rhinotermitidae, 400
mortality,
honey bees, 546
incidents, 546
pesticides, 546
risk assessment, 546
moth,
community structure, 1224
conservation, 1224
spatial structure, 1224
timber harvest, 1224
mountain pine beetle,
4-allylanisole, 871
Dendroctonus ponderosae, 871
estragole, 871
Pinus ponderosa, 871
multiple nutrients,
dietary phosphorus limitation to grasshoppers, 333
ecological stoichiometry, 333
nutritional ecology, 333
nutritional indices, 333
multipredator effect,
Chiapas, 181
intraguild predation, 181
Mexico, 181
resource partitioning, 181
trophic interaction, 181
mutualism, 734, 1471
ant, 1471
biological control, 734
indirect competition, 734
leafhoppers, 1471
scale insects, 734
shade, 1471
shade coffee, 734
shelter, 1471
myrmecophily,
ant-hemipteran mutualism, 143
Coccinellidae, 143
coffee, 143
Mexico, 143
Mythimna separata, 774, 1389
autumn migration, 774
cold, 1389
flight behavior, 774
light trapping, 774
migration behavior, 1389
radar observation, 774
sensitive stage, 1389
starvation, 1389

N

native plant, 478, 808
Apis mellifera, 808
beetle banks, 478
conservation biological control, 478
habitat manipulation, 478
honey bee, 808
insectary plants, 478
invasive species, 808
Triteleia laxa, 808
natural enemies, 121, 1174
agro-ecology, 121
beetle communities, 121
biological control, 121
forest insects, 1174
generalist predators, 121
habitat structure, 1174
invasive species, 1174
Lymantria dispar, 1174
natural enemy,
attractant, 1410
cotton, 1410
plant volatile, 1410
nectar,
anthrone, 996
extrafloral, 996
nutrition, 996
spider, 996
neotenic reproductives,
inbreeding, 883
inclusive fitness, 883
reproductive plasticity, 883
reproductive skew, 883
nest orientation,
insolation, 198
remote sensing, 198
thermoregulation, 198
net reproductive rate, 586, 1538
fecundity, 586
finite rate of increase, 1538
generation time, 1538
instrinsic rate of natural increase, 586
longevity, 586
plant host abundance, 1538
tobacco budworm, 1538
Nilaparvata lugens, 206, 748
Anagrus nilaparvatae, 748
crude fat, 206
host suitability, 748
infestation level, 748
insecticide, 206
reproductive rate, 206
soluble sugar, 206
tritrophic interaction, 748
nitrogen, 575, 1086
cherry fruit, 1086
cherry leaves, 1086
insect, 575
integrated pest management, 575
Rhagoletis indifferens, 1086
sugar, 1086
yield, 575
nonlinear model,
development rate, 348
invasive species, 348
stink bug, 348
nontarget effects,
Aphis gossypii, 1035
Orius sauteri, 1035
transgenic *Bt* cotton, 1035
tri-trophic interactions, 1035
nontarget organisms,
biotechnology, 838
Cry3Bb1, 838
potassium arsenate, 838
protease inhibitor, 838
number of generations,
overwintering, 670
parasitoids, 670
predators, 670
nutrition,
anthrone, 996
extrafloral, 996
nectar, 996
spider, 996
nutritional ecology, 333, 1003
beneficial acclimation hypothesis, 1003
calorimetry, 1003
diet switching, 1003
dietary acclimation hypothesis, 1003
dietary phosphorus limitation to grasshoppers, 333
ecological stoichiometry, 333
multiple nutrients, 333
nutritional indices, 333
nutritional indices,
dietary phosphorus limitation to grasshoppers, 333
ecological stoichiometry, 333
multiple nutrients, 333
nutritional ecology, 333

O

obliquebanded leafroller,
Choristoneura rosaceana, 366
field trapping, 366
flight tunnel, 366

pheromone compounds, 366
odorous house ant,
Argentine ant, 741
competition, 741
Linepithema humile, 741
Tapinoma sessile, 741
Olea europaea L.,
Bactrocera oleae (Gmelin), 764
larval parasitoid, 764
olive grove,
geostatistics, 382
Lobesia botrana, 382
vineyard, 382
operational and environmental factors,
Bemisia tabaci, 1514
insect resistance management, 1514
modeling, 1514
pyriproxyfen, 1514
Oribatid mites,
canopy lichens, 464
distribution patterns, 464
microhabitat specificity, 464
montane forests, 464
Orius sauteri,
Aphis gossypii, 1035
nontarget effects, 1035
transgenic *Bt* cotton, 1035
tri-trophic interactions, 1035
Ostrinia nubilalis,
dispersal, 1280
flight behavior, 1280
flight mill, 1280
migration, 1280
outbreaks,
climate change, 650
Dendroctonus frontalis, 650
Thanasimus dubius, 650
upper lethal temperature, 650
ovarian development,
flight capacity, 374
mating status, 374
overwintering, 670, 1291
Cactoblastis cactorum, 1291
dispersal, 1291
flight, 1291
number of generations, 670
parasitoids, 670
predators, 670
seasonality, 1291
overwintering mortality,
Euproctis chrysorrhoea, 1429
forest defoliator, 1429
invasive species, 1429
population collapse, 1429
oviposition,
fecundity, 520
insect/plant interactions, 520
oviposition preference,
Fraxinus, 230
host range, 230
host selection, 230
invasive pest, 230
oxidative stress,
caterpillar, 1113
phenolic compounds, 1113
superoxide dismutase, 1113
tannin, 1113

P

Panonychus citri,
beneficial arthropods, 555
Dialeurodes citri, 555
Lepidosaphes beckii, 555
Satsuma mandarin, 555
parasitism,
host range, 1231
host-instar utilization, 1231
hyperparasitism, 1231
idiobiont/koinobiont mode, 1231
parasitoid behavior,
Ips pini, 150
microbe interactions, 150
predator behavior, 150
semiochemicals, 150
parasitoid wasp,
courtship, 782
receptivity, 782
signaling, 782
vestigial, 782
parasitoids, 670, 1264, 1307, 1477
Adelges tsugae, 1477
aphid-pathogenic fungi, 1264
bagworm, 1307
Chamaemyiidae, 1477
colonization, 1264
competition, 1307
dissemination, 1264
host discrimination, 1307
host dispersal flight, 1264
host feeding, 1307
number of generations, 670
overwintering, 670
predators, 670, 1477
Tsuga heterophylla, 1477
parental investment,
Euschistus conspersus, 192
mate choice, 192
mating behavior, 192
reproductive behavior, 192
parthenium,
acalypha, 630
development, 630
hibiscus, 630
plumeria, 630
passion vines,
feeding choice tests, 1323
food mixing, 1323
host plant selection, 1323
leaffooted bugs, 1323
pathological effect,
Bombyx mori, 825
Bt rice, 825
pollen grains, 825
toxic effect, 825
pear,
apple, 980
low volume, 980
mating disruption, 980
pest management, 980
pecan,
Beauveria bassiana, 162
biological control, 162
Curculio caryae, 162
pest management, 980, 1136
apple, 980, 1136
competitive attraction, 1136
Cydia pomonella, 1136
low volume, 980
mating disruption, 980, 1136
pear, 980
pesticides,
honey bees, 546
incidents, 546
mortality, 546
risk assessment, 546
pH,
cellulose, 45
development, 45
stable fly, 45
survival, 45
phenolic compounds,
caterpillar, 1113
oxidative stress, 1113
superoxide dismutase, 1113
tannin, 1113
phenology, 472, 696, 914, 1185, 1558
Aphidius ervi, 472
biological control, 1185
Diabrotica virgifera virgifera, 1558
geographical distribution, 472
global warming, 696
host plant, 1185
huanglongbing, 914
life history, 1185
Lixinae, 1185
Pieris rapae, 696
population dynamics, 472
Praon unicum, 472
precipitation, 696
root toughness, 1558
sampling, 914
sticky traps, 914
temperature, 696
trapping, 914
phenology simulation model,
development, 660
Scotinophara lurida, 660
temperature, 660

pheromone compounds,
Choristoneura rosaceana, 366
field trapping, 366
flight tunnel, 366
obliquebanded leafroller, 366
pheromones,
exo-brevicomin, 70
interspecific communication, 70
reproductive isolation, 70
trapping, 70
Phoebe oil,
bark volatiles, 356
Buprestidae, 356
7-*epi*-sesquithujene, 356
Manuka oil, 356
photoperiod,
Anthonomus grandis grandis, 1396
boll weevil, 1396
cotton, 1396
phytophage,
aphids, 1531
resistance, 1531
sprouting, 1531
susceptibility, 1531
Phytoseiidae,
development, 340
life table, 340
pollen, 340
Typhlodromus foenilis, 340
Pierce's disease, 925, 1200
Auchenorrhyncha, 925
egg maturation, 1200
glassy-winged sharpshooter, 925
Hexapoda, 925
invasive species, 1200
leafhopper, 925
Pierce's disease, 925, 1200
Xylella fastidiosa, 1200
Pieris rapae,
global warming, 696
phenology, 696
precipitation, 696
temperature, 696
Pinus ponderosa, 57, 94, 871
4-allylanisole, 871
abundance, 94
degree day, 57
Dendroctonus, 57, 94
Dendroctonus ponderosae, 871
estragole, 871
flight seasonality, 94
Ips, 57, 94
mountain pine beetle, 871
temperature, 57
plant defense theory,
host plant selection, 1548
Japanese beetle, 1548
polyphagy, 1548
Popillia japonica, 1548
plant host abundance,
finite rate of increase, 1538
generation time, 1538
net reproductive rate, 1538
tobacco budworm, 1538
plant pathogen,
Alatae, 1573
growth rate, 1573
host choice, 1573
virus vector, 1573
plant volatile,
attractant, 1410
cotton, 1410
natural enemy, 1410
plant–insect interaction,
compensatory photosynthesis, 1332
defoliation, 1332
folivory, 1332
gas exchange, 1332
plant–insect interaction, 1332
plumeria,
acalypha, 630
development, 630
hibiscus, 630
parthenium, 630
Plutella xylostella,
Brassica oleracea, 38
development rate, 38
Diadegma anurum, 38
temperature, 38
Poa annua,
annual bluegrass weevil, 1438
population fluctuations, 1438
turfgrass, 1438
Podisus maculiventris,
augmentative release, 1241
biological control, 1241
predation dynamics, 1241
predator behavior, 1241
pollen,
development, 340
life table, 340
Phytoseiidae, 340
Typhlodromus foenilis, 340
pollen grains,
Bombyx mori, 825
Bt rice, 825
pathological effect, 825
toxic effect, 825
pollination, 707, 719
bumble bee, 707, 719
habitat management, 707, 719
prairie plants, 707, 719
restoration, 707, 719
polyphagy,
host plant selection, 1548
Japanese beetle, 1548
plant defense theory, 1548
Popillia japonica, 1548
Popillia japonica, 601, 1548
elevated carbon dioxide, 601
fecundity, 601
Glycine max, 601
host plant selection, 1548
Japanese beetle, 1548
longevity, 601
plant defense theory, 1548
polyphagy, 1548
Popillia japonica Newman,
Cotinis nitida L, 608
grapes, 608
green June beetle, 608
Japanese beetle, 608
population collapse,
Euproctis chrysorrhoea, 1429
forest defoliator, 1429
invasive species, 1429
overwintering mortality, 1429
population dynamics, 472, 679, 1043, 1361
Aphidius ervi, 472
Bt cotton, 1043
Choristoneura rosaceana, 679
common cutworm, 1043
Cry1Ac level, 1361
Cydia pomonella, 679
delayed mating, 679
geographical distribution, 472
mating disruption, 679
phenology, 472
Praon unicum, 472
Sesamia inferens, 1361
survival, 1043
transgenic rice lines, 1361
population fluctuations,
annual bluegrass weevil, 1438
Poa annua, 1438
turfgrass, 1438
population genetics,
Diabrotica, 293
DNA markers, 293
microsatellites, 293
western corn rootworm, 293
population structure,
Dendroctonus frontalis, 271
insect pest, 271
microsatellites, 271
southern pine beetle, 271
potassium arsenate,
biotechnology, 838
Cry3Bb1, 838
nontarget organisms, 838
protease inhibitor, 838
prairie grasses,
alternate hosts, 247
Diabrotica barberi, 247
resistance management, 247
prairie plants, 707, 719
bumble bee, 707, 719
habitat management, 707, 719
pollination, 707, 719

restoration, 707, 719
Praon unicum,
Aphidius ervi, 472
geographical distribution, 472
phenology, 472
population dynamics, 472
precipitation, 686, 696
food limitation, 686
global warming, 696
grasshoppers, 686
insect population dynamics, 686
phenology, 696
Pieris rapae, 696
temperature, 696
weather conditions, 686
predation dynamics,
augmentative release, 1241
biological control, 1241
Podisus maculiventris, 1241
predator behavior, 1241
predator behavior, 150, 1241
augmentative release, 1241
biological control, 1241
Ips pini, 150
microbe interactions, 150
parasitoid behavior, 150
Podisus maculiventris, 1241
predation dynamics, 1241
semiochemicals, 150
predator–prey interactions,
enzyme-linked immunosorbent assay, 990
gut content analysis, 990
monoclonal antibody, 990
predator–prey interactions, 990
sampling protocols, 990
predators, 670, 1477
Adelges tsugae, 1477
Chamaemyiidae, 1477
number of generations, 670
overwintering, 670
parasitoids, 670, 1477
Tsuga heterophylla, 1477
preference,
insidious flower bug, 964
prey consumption, 964
soybean aphid, 964
soybean thrips, 964
preference assay,
Aphis glycines, 973
buckthorn, 973
Harmonia axyridis, 973
soybean aphid, 973
preimaginal development,
adult life history, 307
Dacini, 307
host–plant effect, 307
temperature, 307
prey consumption,
insidious flower bug, 964
preference, 964
soybean aphid, 964
soybean thrips, 964
Propylaea japonica,
Aphis gossypii, 29
cotton cultivar, 29
elevated CO_2, 29
tri-trophic interaction, 29
protease inhibitor,
biotechnology, 838
Cry3Bb1, 838
nontarget organisms, 838
potassium arsenate, 838
protein,
feeding behavior, 172
feeding ecology, 172
Rhagoletis indifferens, 172
sugar, 172
Prunus angustifolia,
Prunus persica, 1508
Prunus serotina, 1508
Rosaceae, 1508
Synanthedon pictipes, 1508
Prunus persica,
Prunus angustifolia, 1508
Prunus serotina, 1508
Rosaceae, 1508
Synanthedon pictipes, 1508
Prunus serotina,
Prunus angustifolia, 1508
Prunus persica, 1508
Rosaceae, 1508
Synanthedon pictipes, 1508
Pteromalidae,
Bostrichidae, 1300
Curculionidae, 1300
host finding, 1300
learning, 1300
pupation,
entomopathogenic nematodes, 845
soil moisture, 845
soil texture, 845
putrescine,
cadaverine, 1119
Caribbean fruit fly, 1119
electroantennography, 1119
pyriproxyfen,
Bemisia tabaci, 1514
insect resistance management, 1514
modeling, 1514
operational and environmental factors, 1514

Q

quantile regression,
aerobiology, 87
biological invasions, 87
local indicator of spatial autocorrelation, 87
Lymantria dispar, 87

R

radar observation,
autumn migration, 774
flight behavior, 774
light trapping, 774
Mythimna separata, 774
radiotransmitter,
Anabrus simplex, 889
collective movement, 889
hopper band, 889
katydid, 889
RAPD,
Bacillus thuringiensis, 1354
gene flow, 1354
Mediterranean corn borer, 1354
transgenic maize, 1354
receptivity,
courtship, 782
parasitoid wasp, 782
signaling, 782
vestigial, 782
redbay,
ambrosia beetle, 442
climate matching, 442
cost-weighted distance, 442
Xyleborus glabratus, 442
regulated water deficit,
choice tests, 787
glassy-winged sharpshooter, 787
Xylella fastidiosa, 787
relative humidity,
degree-days, 568
larval development, 568
life table, 568
management, 568
remote sensing,
insolation, 198
nest orientation, 198
thermoregulation, 198
repellency,
Agriotes obscurus, 534
dusky wireworm, 534
insecticide, 534
reproduction,
Heterorhabditis bacteriophora, 1272
Steinernema feltiae, 1272
thermal adaptation, 1272
virulence, 1272
reproductive behavior,
Euschistus conspersus, 192
mate choice, 192
mating behavior, 192
parental investment, 192

reproductive incompatibility,
Conotrachelus nenuphar, 1208
interstrain matings, 1208
reproductive isolation,
exo-brevicomin, 70
interspecific communication, 70
pheromones, 70
trapping, 70
reproductive plasticity,
inbreeding, 883
inclusive fitness, 883
neotenic reproductives, 883
reproductive skew, 883
reproductive rate,
crude fat, 206
insecticide, 206
Nilaparvata lugens, 206
soluble sugar, 206
reproductive skew,
inbreeding, 883
inclusive fitness, 883
neotenic reproductives, 883
reproductive plasticity, 883
resistance,
aphids, 1531
phytophage, 1531
sprouting, 1531
susceptibility, 1531
resistance management,
alternate hosts, 247
Diabrotica barberi, 247
prairie grasses, 247
resource partitioning,
Chiapas, 181
intraguild predation, 181
Mexico, 181
multipredator effect, 181
trophic interaction, 181
restoration, 707, 719
bumble bee, 707, 719
habitat management, 707, 719
pollination, 707, 719
prairie plants, 707, 719
Rhagoletis indifferens, 172, 1086
cherry fruit, 1086
cherry leaves, 1086
feeding behavior, 172
feeding ecology, 172
nitrogen, 1086
protein, 172
sugar, 172, 1086
Rhagoletis pomonella,
apple, 1154
diapause, 1154
Rhinotermitidae,
breeding system, 400
Isoptera, 400
microsatellite genotyping, 400
morphometry, 400
rice,
compensation, 796
Diatraea saccharalis, 796
injury, 796
sugarcane borer, 796
risk assessment, 1, 546
Bacillus thuringiensis, 1
experimental design, 1
genetically modified, 1
honey bees, 546
incidents, 546
mortality, 546
pesticides, 546
root toughness,
Diabrotica virgifera virgifera, 1558
phenology, 1558
Rosaceae,
Prunus angustifolia, 1508
Prunus persica, 1508
Prunus serotina, 1508
Synanthedon pictipes, 1508

S

sample completeness,
butterflies, 1313
day-flying moths, 1313
estimating species richness, 1313
sampling fraction, 1313
sampling,
huanglongbing, 914
phenology, 914
sticky traps, 914
trapping, 914
sampling fraction,
butterflies, 1313
day-flying moths, 1313
estimating species richness, 1313
sample completeness, 1313
sampling protocols,
enzyme-linked immunosorbent assay, 990
gut content analysis, 990
monoclonal antibody, 990
predator–prey interactions, 990
saponins,
alfalfa, 947
arthropods, 947
chemical defense, 947
diversity cascades, 947
Satsuma mandarin,
beneficial arthropods, 555
Dialeurodes citri, 555
Lepidosaphes beckii, 555
Panonychus citri, 555
scale insects,
biological control, 734
indirect competition, 734
mutualism, 734
shade coffee, 734
Scarabaeinae,
African savannas, 431
bait types, 431
Botswana, 431
trophic associations, 431
Schinus terebinthifolius,
ArcGIS mapping, 862
biological control, 862
developmental rate, 862
Episimus utilis, 862
Schizaphis graminum,
Lysiphlebus testaceipes, 1063
supercooling point, 1063
winter hardiness, 1063
Scotinophara lurida,
development, 660
phenology simulation model, 660
temperature, 660
searching speed,
Diaprepes root weevil, 1565
egg parasitoid, 1565
host plants, 1565
trichomes, 1565
seasonal development,
cole crops, 907
swede midge, 907
seasonal dynamics,
behavior, 1373
cotton gin trash, 1373
soil moisture preference, 1373
seasonality,
Cactoblastis cactorum, 1291
dispersal, 1291
flight, 1291
overwintering, 1291
secondary production,
Chironomidae, 1162
sex dimorphism, 1162
sex ratios, 1162
tropical stream, 1162
seed bank,
Centaurea stoebe, 510
Larinus spp., 510
seed production, 510
Urophora affinis, 510
seed production,
Centaurea stoebe, 510
Larinus spp., 510
seed bank, 510
Urophora affinis, 510
semiochemicals,
Ips pini, 150
microbe interactions, 150
parasitoid behavior, 150
predator behavior, 150
sensitive stage,
cold, 1389
migration behavior, 1389
Mythimna separata, 1389

starvation, 1389
Sesamia inferens,
Cry1Ac level, 1361
population dynamics, 1361
transgenic rice lines, 1361
sex dimorphism,
Chironomidae, 1162
secondary production, 1162
sex ratios, 1162
tropical stream, 1162
sex pheromone, 1126, 1416
Anomala orientalis, 1126
dynamic head space sampling, 1416
electroantennogram, 1416
mating disruption, 1126
tea mosquito, 1416
turfgrass, 1126
wind tunnel, 1416
(Z) and (E)-7-tetradecen-2-one formulation, 1126
sex ratios,
Chironomidae, 1162
secondary production, 1162
sex dimorphism, 1162
tropical stream, 1162
shade,
ant, 1471
leafhoppers, 1471
mutualism, 1471
shelter, 1471
shade coffee,
biological control, 734
indirect competition, 734
mutualism, 734
scale insects, 734
shelter,
ant, 1471
leafhoppers, 1471
mutualism, 1471
shade, 1471
signaling,
courtship, 782
parasitoid wasp, 782
receptivity, 782
vestigial, 782
sika deer,
ecosystem engineering, 938
habitat fragmentation, 938
vegetation structure, 938
silver birch,
chitinase, 1582
ecological interactions, 1582
herbivory, 1582
transgenic tree, 1582
Sitona discoideus,
developmental rate, 1381
embryo development, 1381
modeling, 1381
temperature, 1381
soil moisture,
entomopathogenic nematodes, 845
pupation, 845
soil texture, 845
soil moisture preference,
behavior, 1373
cotton gin trash, 1373
seasonal dynamics, 1373
soil temperature,
animal foraging, 857
daily variations, 857
harvester ants, 857
trail system, 857
soil texture,
entomopathogenic nematodes, 845
pupation, 845
soil moisture, 845
Solenopsis,
Black Belt Prairie, 453
ecological assembly, 453
Formicidae, 453
Mississippi, 453
Solenopsis invicta,
Antonina graminis, 487
Cynodon dactylon, 487
honeydew, 487
invasive species, 487
soluble sugar,
crude fat, 206
insecticide, 206
Nilaparvata lugens, 206
reproductive rate, 206
southern pine beetle,
Dendroctonus frontalis, 271
insect pest, 271
microsatellites, 271
population structure, 271
Soxhlet extraction,
Dendroctonus pseudotsugae, 850
Douglas-fir beetle, 850
fatreserves, 850
insect flight, 850
soybean aphid, 964, 973, 1258
ant attendance, 1258
Aphis glycines, 973
buckthorn, 973
Coccinellidae, 1258
Harmonia axyridis, 973
insidious flower bug, 964
Monomorium, 1258
preference, 964
preference assay, 973
prey consumption, 964
soybean thrips, 964
soybean thrips,
insidious flower bug, 964
preference, 964
prey consumption, 964
soybean aphid, 964
spatial dynamics,
cabbage root fly, 409
crop rotation, 409
Delia radicum, 409
spatial ecology, 409
spatial ecology,
cabbage root fly, 409
crop rotation, 409
Delia radicum, 409
spatial dynamics, 409
spatial structure,
community structure, 1224
conservation, 1224
moth, 1224
timber harvest, 1224
species richness,
beta diversity, 1488
Calathus, 1488
completeness of inventories, 1488
Iberian Peninsula, 1488
spider,
anthrone, 996
extrafloral, 996
nectar, 996
nutrition, 996
spider diversity,
Argentina, 419
contiguous habitats, 419
Humid Chaco ecoregion, 419
Mburucuya National Park, 419
spinosad,
Chrysoperla externa, 1252
cypermethrin, 1252
endosulfan, 1252
methoxyfenozide, 1252
sprouting,
aphids, 1531
phytophage, 1531
resistance, 1531
susceptibility, 1531
spruce beetle,
classification and regression tree analysis, 281
climate, 281
disturbance ecology, 281
Engelmann spruce, 281
stable fly,
cellulose, 45
development, 45
pH, 45
survival, 45
starvation,
cold, 1389
migration behavior, 1389
Mythimna separata, 1389
sensitive stage, 1389
Steinernema feltiae,
Heterorhabditis bacteriophora, 1272
reproduction, 1272
thermal adaptation, 1272

virulence, 1272
sticky traps,
huanglongbing, 914
phenology, 914
sampling, 914
trapping, 914
stink bug,
development rate, 348
invasive species, 348
nonlinear model, 348
Stomoxys,
adult feeding, 625
blood, 625
sugar, 625
stored product, 131, 1105
Acanthoscelides obtectus, 131
biological control, 131
Callosobruchus spp., 131
development, 1105
life table, 1105
Liposcelis badia, 1105
temperature, 1105
Zabrotes subfasciatus, 131
Stratiomyidae,
black soldier fly larvae, 1525
dairy manure, 1525
Escherichia coli, 1525
Hermetia illucens, 1525
subterranean termites,
fungal pathogen, 755
fungal repellence, 755
integrated pest management, 755
landscape mulch management, 755
sugar, 172, 625, 1086
adult feeding, 625
blood, 625
cherry fruit, 1086
cherry leaves, 1086
feeding behavior, 172
feeding ecology, 172
nitrogen, 1086
protein, 172
Rhagoletis indifferens, 172, 1086
Stomoxys, 625
sugarcane borer,
compensation, 796
Diatraea saccharalis, 796
injury, 796
rice, 796
sunflower beetle,
arbuscular mycorrhizal fungi, 1017
Helianthus annuus, 1017
trait-mediated indirect effects, 1017
Zygogramma exclamationis, 1017
supercooling point, 1063, 1074
body size, 1074, 1074
cooling rate, 1074
cooling rate interaction, 1074
fire ant, 1074
Lysiphlebus testaceipes, 1063
Schizaphis graminum, 1063
winter hardiness, 1063
superoxide dismutase,
caterpillar, 1113
oxidative stress, 1113
phenolic compounds, 1113
tannin, 1113
survival, 16, 45, 1043, 1145
Bt cotton, 1043
capture-recapture, 1145
cellulose, 45
common cutworm, 1043
Cormack-Jolly-Seber, 1145
development, 45
developmental time, 16
Helicoverpa armigera, 16
life table, 16
pH, 45
population dynamics, 1043
stable fly, 45
temperature, 16
vagility, 1145
susceptibility,
aphids, 1531
phytophage, 1531
resistance, 1531
sprouting, 1531
swede midge,
cole crops, 907
seasonal development, 907
sweetpotato whitefly,
Bemisia tabaci biotype B, 391
biological control, 391
Encarsia spp., 391
Eretmocerus spp., 391
symbiont,
emerald ash borer, 1344
forest insect, 1344
insect gut, 1344
invasive species, 1344
symbiosis,
bark beetles, 956
Histiogaster, 956
Hyphomycetes, 956
Leptographium, 956
Synanthedon pictipes,
Prunus angustifolia, 1508
Prunus persica, 1508
Prunus serotina, 1508
Rosaceae, 1508
synchrony,
Adelges tsugae, 1498
classical biological control, 1498
exclusion cage, 1498
field insectary, 1498

T

tannin,
caterpillar, 1113
oxidative stress, 1113
phenolic compounds, 1113
superoxide dismutase, 1113
Tapinoma sessile,
Argentine ant, 741
competition, 741
Linepithema humile, 741
odorous house ant, 741
tea mosquito,
dynamic head space sampling, 1416
electroantennogram, 1416
sex pheromone, 1416
wind tunnel, 1416
temperature, 16, 38, 57, 307, 660, 696, 1105, 1381
adult life history, 307
Brassica oleracea, 38
Dacini, 307
degree day, 57
Dendroctonus, 57
development, 660, 1105
developmental rate, 38, 1381
developmental time, 16
Diadegma anurum, 38
embryo development, 1381
global warming, 696
Helicoverpa armigera, 16
host–plant effect, 307
Ips, 57
life table, 16, 1105
Liposcelis badia, 1105
modeling, 1381
phenology, 696
phenology simulation model, 660
Pieris rapae, 696
Pinus ponderosa, 57
Plutella xylostella, 38
precipitation, 696
preimaginal development, 307
Scotinophara lurida, 660
Sitona discoideus, 1381
stored product, 1105
survival, 16
Tephritidae,
Anastrepha obliqua, 876
calling behavior, 876
Diptera, 876
male volatiles, 876
Thanasimus dubius,
climate change, 650
Dendroctonus frontalis, 650
outbreaks, 650
upper lethal temperature, 650
thelytoky,
arrhenotoky, 1422
haplodiploidy, 1422
Thysanoptera, 1422

Wolbachia, 1422
thermal adaptation,
Heterorhabditis bacteriophora, 1272
reproduction, 1272
Steinernema feltiae, 1272
virulence, 1272
thermoregulation,
insolation, 198
nest orientation, 198
remote sensing, 198
Thysanoptera,
arrhenotoky, 1422
haplodiploidy, 1422
thelytoky, 1422
Wolbachia, 1422
timber harvest,
community structure, 1224
conservation, 1224
moth, 1224
spatial structure, 1224
tobacco budworm, 1081, 1538
cottonseed, 1081
finite rate of increase, 1538
generation time, 1538
Gossypium hirsutum, 1081
(+)-gossypol, 1081
net reproductive rate, 1538
plant host abundance, 1538
tomato,
electrical penetration graph, 213
Frankliniella fusca, *213*
Frankliniella occidentalis, 213
tomato spotted wilt virus,
epidemiology, 79
insect dispersal, 79
toxic effect,
Bombyx mori, 825
Bt rice, 825
pathological effect, 825
pollen grains, 825
trade-off,
body size, 51
encapsulation, 51
habitat quality, 51
immunity, 51
trail system,
animal foraging, 857
daily variations, 857
harvester ants, 857
soil temperature, 857
trait-mediated indirect effects,
arbuscular mycorrhizal fungi, 1017
Helianthus annuus, 1017
sunflower beetle, 1017
Zygogramma exclamationis, 1017
transgenic *Bt* cotton, 263, 1035
Aphis gossypii, 263, 1035
Chrysopa pallens, 263
nontarget effects, 1035
Orius sauteri, 1035
tri-trophic interactions, 263, 1035
transgenic corn,
Bt accumulation in soil, 255
Cry1F protein, 255
environmental risk, 255
enzyme-linked immunosorbent assay, 255
transgenic maize,
Bacillus thuringiensis, 1354
gene flow, 1354
Mediterranean corn borer, 1354
RAPD, 1354
transgenic rice lines,
Cry1Ac level, 1361
population dynamics, 1361
Sesamia inferens, 1361
transgenic tree,
chitinase, 1582
ecological interactions, 1582
herbivory, 1582
silver birch, 1582
trapping, 70, 914
exo-brevicomin, 70
huanglongbing, 914
interspecific communication, 70
phenology, 914
pheromones, 70
reproductive isolation, 70
sampling, 914
sticky traps, 914
treatments,
biological management, 505
Diptera, 505
entomopathogenic bacterium, 505
microbial insecticide, 505
tri-trophic interaction, 29, 263, 1035
Aphis gossypii, 29, 263, 1035
Chrysopa pallens, 263
cotton cultivar, 29
elevated CO_2, 29
nontarget effects, 1035
Orius sauteri, 1035
Propylaea japonica, 29
transgenic *Bt* cotton, 263, 1035
trichomes,
Diaprepes root weevil, 1565
egg parasitoid, 1565
host plants, 1565
searching speed, 1565
Triteleia laxa,
Apis mellifera, 808
honey bee, 808
invasive species, 808
native plant, 808
tritrophic interaction, 748, 1025
Ageniaspis citricola, 1025
Anagrus nilaparvatae, 748
Cirrospilus neotropicus, 1025
citrus leafminer, 1025
host selection, 1025
host suitability, 748
infestation level, 748
Nilaparvata lugens, 748
trophic associations,
African savannas, 431
bait types, 431
Botswana, 431
Scarabaeinae, 431
trophic interaction,
Chiapas, 181
intraguild predation, 181
Mexico, 181
multipredator effect, 181
resource partitioning, 181
trophic selection,
Cassidinae, 224
feeding niches, 224
Hispinae, 224
Zingiberales, 224
tropical stream,
Chironomidae, 1162
secondary production, 1162
sex dimorphism, 1162
sex ratios, 1162
Tsuga heterophylla, 494, 1477
Adelges tsugae, 494, 1477
Chamaemyiidae, 1477
Laricobius nigrinus, 494
Leucopis argenticollis, 494
Leucopis atrifacies, 494
parasitoids, 1477
predators, 1477
turfgrass, 1126, 1438
annual bluegrass weevil, 1438
Anomala orientalis, 1126
mating disruption, 1126
Poa annua, 1438
population fluctuations, 1438
sex pheromone, 1126
(Z) and (E)-7-tetradecen-2-one formulation, 1126
Typhlodromus foenilis,
development, 340
life table, 340
Phytoseiidae, 340
pollen, 340

U

upper lethal temperature,
climate change, 650
Dendroctonus frontalis, 650
outbreaks, 650
Thanasimus dubius, 650
Urophora affinis,
Centaurea stoebe, 510
Larinus spp., 510

seed bank, 510
seed production, 510
Utetes anastrephae,
Anastrepha obliqua, 110
Anastrepha suspensa, 110
Mangifera indica, 110

V

vagility,
capture-recapture, 1145
Cormack-Jolly-Seber, 1145
survival, 1145
vector performance,
alternate weed hosts, 592
insect vectors, 592
vector reservoir, 592
viral inoculum source, 592
vector reservoir,
alternate weed hosts, 592
insect vectors, 592
vector performance, 592
viral inoculum source, 592
vegetation structure,
ecosystem engineering, 938
habitat fragmentation, 938
sika deer, 938
vestigial,
courtship, 782
parasitoid wasp, 782
receptivity, 782
signaling, 782
vineyard,
geostatistics, 382
Lobesia botrana, 382
olive grove, 382
viral inoculum source,
alternate weed hosts, 592
insect vectors, 592
vector performance, 592
vector reservoir, 592
virulence,
Heterorhabditis bacteriophora, 1272
reproduction, 1272
Steinernema feltiae, 1272
thermal adaptation, 1272
virus vector,
Alatae, 1573
growth rate, 1573
host choice, 1573
plant pathogen, 1573

W

weather conditions,
food limitation, 686
grasshoppers, 686
insect population dynamics, 686
precipitation, 686
western corn rootworm,
Diabrotica, 293
DNA markers, 293
microsatellites, 293
population genetics, 293
wind tunnel,
dynamic head space sampling, 1416
electroantennogram, 1416
sex pheromone, 1416
tea mosquito, 1416
winter hardiness,
Lysiphlebus testaceipes, 1063
Schizaphis graminum, 1063
supercooling point, 1063
Wolbachia,
arrhenotoky, 1422
haplodiploidy, 1422
thelytoky, 1422
Thysanoptera, 1422
wood borers,
Agrilus planipennis, 242
coevolution, 242
host plant resistance, 242
invasive species, 242

X

Xyleborus glabratus,
ambrosia beetle, 442
climate matching, 442
cost-weighted distance, 442
redbay, 442
Xylella fastidiosa, 787, 1049, 1200
behavior, 1049
choice tests, 787
egg maturation, 1200
glassy-winged sharpshooter, 787, 1049
invasive species, 1200
management, 1049
Pierce's disease, 1200
regulated water deficit, 787

Y

yield,
insect, 575
integrated pest management, 575
nitrogen, 575

Z

(Z) and (E)-7-tetradecen-2-one formulation,
Anomala orientalis, 1126
mating disruption, 1126
sex pheromone, 1126
turfgrass, 1126
Zabrotes subfasciatus,
Acanthoscelides obtectus, 131
biological control, 131
Callosobruchus spp., 131
stored products, 131
Zingiberales,
Cassidinae, 224
feeding niches, 224
Hispinae, 224
trophic selection, 224
zoophytophagy,
Coleomegilla maculata, 525
Cry1Ab endotoxin, 525
dietary mixing, 525
Harmonia axyridis, 525
Zygogramma exclamationis,
arbuscular mycorrhizal fungi, 1017
Helianthus annuus, 1017
sunflower beetle, 1017
trait-mediated indirect effects, 1017